GAJAH

Number 24
2006
ISSN: 1391-1996

Journal of the Asian Elephant Specialist Group



GAJAH

Jounal of the Asian Elephant Specialist Group
Number 24: (2006)
ISSN: 1391-1996

GAJAH is the official journal of the Asian Elephant Specialist Group (AsESG) of the Species Survival Commission (SSC) of the World Conservation Union (IUCN). The journal is intended as a medium for communication by members of the AsESG of important issues that concern the conservation and management of the Asian Elephant (*Elephas maximus*) both in the wild and in captivity. GAJAH welcomes communications and research papers on all aspects of the Asian elephant. GAJAH is aimed at professionals, biologists and academics carrying out research on Asian elephant, government and non-government organizations involved in its conservation, and interested members of the general public. All articles published in GAJAH are deemed to reflect the individual views of the authors and not the official points of view, either of the Asian Elephant Specialist Group (AsESG) or the Species Survival Commission (SSC). GAJAH is a non-profit publication that is supported by financial assistance from the U.S. Fish and Wildlife Service.

Editor

Charles Santiapillai
Department of Zoology
University of Peradeniya
Sri Lanka
e-mail: csantiapillai@gmail.com

Editorial Board

Richard F. W. Barnes
Department of Biology 0116
University of California at San Diego
La Jolla, CA 92093-0116
USA
e-mail: rbarnes@uscd.edu

Jayantha Jayewardene
Biodiversity & Elephant Conservation Trust
615/32, Rajagiriya Gardens
Nawala Road, Rajagiriya
Sri Lanka
e-mail: romalijj@eureka.lk

M. Philip Kahl
100 Mountain Road
Pine Valley
Sedona, Arizona 86351
USA
e-mail: musthbull@sedona.net

D.K. Lahiri-Choudhury
45 Suhasini Ganguly Sarani
Calcutta 700 025
India
e-mail: dklc@cal.vsnl.net.in

Heidi S. Riddle
Riddle's Elephant and Wildlife Sanctuary,
Post Office Box 715,
Greenbrier,
Arkansas 72058, USA
e-mail: gajah@alltel.net

Raman Sukumar
Centre for Ecological Sciences
Indian Institute of Science
Bangalore
India
e-mail: rsuku@ces.iisc.ernet.in

GAJAH

Journal of the Asian Elephant Specialist Group
Volume 24: (2006)
ISSN: 1391-1996

The production of GAJAH is financed by the U.S. Fish and Wildlife Service through IUCN - The World Conservation Union and the World Wide Fund for Nature (WWF)

IUCN
The World Conservation Union

WWF

Cover: "Seblat Elephant Conservation Center and Conservation Response Unit Staff members, Bengkulu Province, Sumatra".

Photo: Heidi Riddle for International Elephant Foundation (IEF)

Layouts & Cover Designed by Rangana Wickramathilaka
Printed at Digital TELEPRINTS (Pvt) Ltd - Kandy, Sri Lanka.

GAJAH

Journal of the Asian Elephant Specialist Group (Number 24, 2006)

Contents

Conservation News 1
Comment 3
Charles Santiapillai

Papers

Genetic assessment of Bornean elephants: origin and conservation implications
Prithiviraj Fernando, John Payne, Geoffrey Davison, Raymond J. Alfred, Michael Stuewe and Don J. Melnick 5

Establishing a sustainable model for the long-term conservation of the elephant in Sri Lanka
Ravi Corea 13

Survey of Elephants in the Mannar District, Sr Lanka
S. Wijeyamohan, S.R.B. Dissanayake and Charles Santiapillai 19

Mahouts and their elephants working as Conservation Response Unit in Sumatra
Wahdi Azmi, Heidi S. Riddle, Agung Setyabudi, Wiranto, Andi Basrul, Edy Sunardi, Diding M. Ichsan and Selamat Hardi 35

The shifting chemical signals of Musth
Heidi S. Riddle, David R. Greenwood and L.E.L. (Bets) Rasmussen 39

Lipid profile of captive Sri Lankan elephants
W.D. Ratnasooriya, G.K.H. de Alwis, R.D. Wijesekera, R.M. Amarasinghe and D. Perera 45

Short Communications

Information on elephant keepers in Sri Lanka
A. Dangolla and I.D. Silva 51

Elephant Endotheliotropic Herpes Virus (EEHV)
Charlie Gray 55

Historical Notes

History and culture of elephants in Sri Lanka
Jayantha Jayewardene 57

Book Review

Endangered Elephants: Past, Present and Future 61

Miscellaneous

IUCN/SC Asia Elephant Specialist Group – Announcement of new Co-Chairs
Ajay Desai and Simon Hedges 63

Ringling Bros. and Barnum & Bailey Commits $ 180,000 to Smithsonian's National Zoo to Continue Critical Elephant Herpes Research 65

International Elephant Foundation 2007 Elephant Conservation and Research Funding Support
Deborah Olson 67

Conservation News

Sri Lanka

Re-introduction of the insurance scheme for mahouts

The Diyawadana Nilame of the Sacred Temple of the Tooth (The Dalada Maligawa, Kandy), Neelanga Dela Bandara plans to re-introduce the insurance scheme for mahouts following an unfortunate incident of a mahout named Mr. M. Piyasena killed by his elephant while he attempted to give it a wash He was 59-year old father of four children. According to an eye witness, the elephant concerned was boisterous for a while forcing Piyasena to use the goad to control it. The elephant became so furious that he took the goad and broke it in two, before turning on the mahout. This was the second incident of elephants in captivity killing their mahouts within two months. In the past two years, more than six mahouts have been killed by enraged elephants in Kandy alone.

Source: The Sunday Times (Sri Lanka) November 6, (2005).

USA

Sanctuary receives Excellence Award

Riddle's Elephant & Wildlife Sanctuary has become the first recipient of the Institution of Excellence Award given by the Elephant Managers Association (EMA), which is a non-profit organization of professional elephant handlers, administrators, veterinarians, researchers and elephant enthusiasts. The award was handed out in the Fall of 2005 during the EMA annual conference hosted by the Portand Zoo in Oregon. The award was granted "for contributions to elephant training, research and care performed with professionalism, integrity and passion". The Sanctuary is operated by an Arkansas non-profit organization dedicated to preserving both Asia and Africa elephants. It offers permanent refuge to any elephant in need. The facility provides training and education in elephant management and conservation and conducts scientific study in the health and physiology of elephants.

Source: Log Cabin Democrat, December 24, (2005).

Workshop on Ultrasound and Veterinary Procedures

The 9th Ultrasound and Veterinary Procedures Workshop for Wildlife Veterinarians was hosted by the Riddle's Elephant & Wildlife Sanctuary in May 2005 at Greenbrier, Arkansas. World renowned elephant veterinarian Dr. Dennis Schmitt, assisted by the sanctuary staff, taught the course. Veterinarians from international elephant facilities (Canada, USA and Sri Lanka) attended the course and shared experiences.

Source: Pachyderm Periodical, 15: 3 (2005).

Elephant program at the Smithsonian Institution's National Zoological Park

Elephants in North American Zoos number approximately less than 275 among 70 or so widely dispersed cities. AZA institutions attract over 150 million visitors, more than all professional sports combined. American Zoos have a unique ability to focus and educate these visitors on the critical conservation issues of this century. Within North America itself, there is tremendous need to establish breeding bull groups as management strategies to dramatically increase the captive population and improve genetic diversity. Captive elephants are not breeding fast enough to keep up with natural mortality. The Smithsonian Institution National Zoological Park can change all this now and make a critical leadership step. It plans to proceed with an ambitious plan to renovate its present elephant exhibit. The new facility will take up an 8-acre parcel of land at the Zoo, with approximately 3 acres of ground space for a group comprising 8 Asian elephants. In addition, the Zoo plans to expand its capability to conduct Asian elephant research and propagation by eventually developing a facility at its Front Royal campus – a 3,200 acre parcel of land that constitutes the Zoo's Conservation & Research Center.

Source: Smithsonian Institution National Zoological Park (2006).

Cambodia

Community participation in elephant conservation in Cambodia

As wild elephant habitat is eroded by the spread of cultivation and over exploitation of forest resources, wild elephants resort to feeding on crops. In this context, the resolution of human-elephant conflict to prevent death or injury on both sides, is no less important than the conservation of elephant habitat. A recent survey of Cambodian rural communities indicates that the work of Fauna & Flora International (FFI)'s Human-Elephant Conflict (HEC) teams is already changing attitudes to Asian elephants. Community leaders interviewed by Tuy Sureivathana, one of FFI's local counterparts in Cambodia's Ministry of Environment, revealed that villagers no longer retaliate against elephants that raid their crops or damage their property. FFI team has been instrumental in dissuading people from using spring traps and even poison against the elephants. The mitigation of the HEC is a vital element of FFI's Asian Elephant Conservation Programme, which also works to prevent poaching, protect habitats and gather crucial scientific data.

Source: FFI Update 2 (2005).

Kenya

Elephant translocation

The elephant population in the Tsavo National Park in the 1960s numbered 40,000. However, poaching for ivory has since reduced the numbers to less than 7,000. In August 2005, the Kenya Wildlife Service (KWS) began rounding up 400 of the 700 elephants in Shimba Hills and moving them to Tsavo East, a much bigger park in southern Kenya with the hope of repopulating the national park. Family groups were selected for translocation by marksmen in helicopters who fired tranquilizer darts at their targets, before loading the elephants into a giant steel crate for the 300 km journey. KWS hopes to track the relocated elephants using GPS collars, fitted to the matriarchs who led each family. The operation will cost the KWS about $ 3 million, and is welcomed by the villagers near Shimba Hills who complain that elephants regularly stray from the park and raid their crops. A previous attempt in 1996 to translocate about 30 bull elephants from Shimba Hills to parkland about 65 km away resulted in the animals becoming disoriented and trekking back towards the coast, eventually reaching another forest reserve further north. Conservationists are divided in their views about such massive translocations of elephants. Daphne Sheldrick who runs the David Sheldrick Wildlife Trust feels that the money could be better spent in erecting a proper barrier fence. Besides, moving elephants from such lush habitats in Shimba Hills to the dry area of Tsavo East national park, may be harmful. But KWS is of the view that unless the overcrowding is eased, elephants will continue to break down barriers.

Source: Jeevan Vasagar in Guardian Weekly, September 2-8, (2005).

South Africa, Mozambique & Zimbabwe

World's biggest Transfrontier Park

An initiative of the Peace Parks Foundation (PPF), the proposed Great Limpopo Transfrontier Park will be the world's biggest animal kingdom covering an area equivalent to half of Scotland and crossing the borders of South Africa, Mozambique and Zimbabwe. These countries are merging three game reserves, thereby creating a 35,000 sq. km conservation area that will expand eventually to cover 100,000 sq. km. The project has been praised as an example of regional cooperation and sustainable development, raising foreign investment and creating much-needed jobs. But some people are not happy. In Mozambique, people from Salani village fear that they will no longer be the hunters but the hunted. Some villagers complain that animal rights have taken precedence over human rights, while others are waiting to see what happens.

Source: Kristy Siegfried in Guardian Weekly, Oct. 28-Nov.3, (2005).

Malaysia

Asian Elephant Range States Meeting

The Meeting of the Asian Elephant Range States was held on January 24-26, 2006 in Malaysia. The meeting commenced with an inaugural session, where Dr. Holly Dublin - the Chair of IUCN Species Survival Commission delivered the key note address, followed by addresses made by the Director General of the Department of Wildlife and National Parks, Malaysia (Perhilitan), and the Parliamentary Secretary of the Ministry of Natural Resources and Environment (NRE). Dr. Dublin highlighted the importance of this meeting, considering the participation of all 13 range states of the Asian Elephant at this occasion to discuss conservation issues relating to the Asian Elephant. The DG of Perhilitan highlighted several issues related to the conservation of Asian Elephants, and stressed the need for dialogue and collaboration between range states. Subsequent to the inaugural session, Dr. Dublin gave a general introduction to the meeting, stressing that the objective was to initiate a dialogue among range states to discuss issues and experiences, and develop consensus towards achieving long-term conservation of Asian Elephants. Dr. Meenakshi Nagendran from the US Fish and Wildlife Service (USFWS) Asian Elephant Conservation Fund highlighted the participation of all 13 range states at this meeting as a great achievement and stated that this was a good opportunity for the range states to discuss the whole range of issues related to the conservation of Asian Elephants, and she indicated that the USFWS would be able to help implement some of the follow-up actions suggested at this meeting through their grant programme.

The key issues highlighted were (a) population management, such as addressing cross border problems, is hindered by information gaps on distribution and population status; (b) lack of standardized methods for population survey and monitoring leading to a lack of good quality data about population distribution and status; and (c) dangers of designing conservation strategies based on guesses about population sizes and trends (efficient allocation of resources for conservation of Asian elephants requires much better data). The recommendations included the need to establish a standardized database on the status and distribution of the Asian Elephant; and the update of existing information through well-designed field surveys using modern peer-reviewed techniques.

Source: AsESG Co-Chairs (2006).

Comment

In a recent issue of the journal *Nature* (438: 1097-1098; December 22-29, 2005), David R. Greenwood, Dan Comeskey, Martin B. Hunt & L. Elizabeth L. Rasmussen have published a paper that deals with the impact of the pheromones associated with the phenomenon of musth on reproductive behaviour of Asian elephants. The word 'musth' means 'intoxicated' in Urdu, and the phenomenon has been known to elephant handlers and keepers for many centuries in India and Sri Lanka. But it is only now, thanks to the studies of Dr. Bets Rasmussen and her colleagues (both in the USA and New Zealand) we seem to understand the role played by the pheromones involved in musth. During musth, Asian bull elephants go through a heightened period of sexual and aggressive activity. The hormone testosterone controls such sexual and aggressive behaviour in bull elephants. In Africa, a musth bull walks with an assertive, purposeful stride and holds the head higher than the shoulders most of the time (Poole, 1987). Kahl & Armstrong (2002) refer to such musth walk as the "John Wayne walk" 'because of its self-confident, take-charge swagger'.

Musth is exhibited only by mature males, usually over 24 years of age (Poole, 1987) and the behaviour of the bulls in musth is the same in both Asian and African elephants (Kahl & Armstrong, 2002). Musth in Asian elephants is an annual phenomenon, and its duration varies from individual to individual depending on age, physical condition and social status. It may last from a few weeks to even months. In captive situation, bulls in musth are always separated from conspecifics and cared for until the musth period is over. When elephants get out of musth, they tend to become more social, amicable and manageable. Although non-musth bull elephants can mate successfully, musth improves the competitive ability of the bulls in their efforts to have access to estrus females. In Asia, older bulls in musth seem to have a significant reproductive advantage over younger bulls. In the captive elephant facility known as the Elephant Orphanage at Pinnawala, Sri Lanka, almost all the young ones that were born were sired by just a few old bulls.

During musth, Asian bull elephants secrete a powerful, pungent smelling fluid from the temporal glands situated on the face. A previous study carried out by Rasmussen *et al.* (2002) on musth in Asian elephants has shown that young socially immature bulls release honey-like odours to avoid conflict with mature bulls, while older bulls in musth broadcast foul-smelling odours to deter young bulls. This was a significant finding that confirmed what the ancient Hindu poetry refers to the phenomenon of bees being attracted by the secretions of the temporal glands in young musth elephants.

As Greenwood *et al.* (2005) point out, musth among Asian elephants is mediated by the release of a pheromone called frontalin, which exists in two chiral forms or molecular mirror images or enantiomers. They are referred to as + and – enantiomers. The researchers have found that these enantiomers of frontalin are released in specific ratio that depends on the animal's state of musth and age. Depending on the ratio of the enantiomers released, other bulls and cows in a population may react and respond in different manners.

Asian bull elephants reach sexual maturity when they are between 12-15 years old, but until they become socially mature, they have very little reproductive success in the wild. Social maturity may take another 5-8 years. The study also shows that frontalin is released in young bulls in late teens and the secretion increases 15-fold in 25 year age span – the time when the bulls become socially mature. As young bulls become sexually mature, they secrete more of the + variety of frontalin than the – variety but as they become socially mature, the proportions of the two enantiomers become equal (1:1 ratio). It is this ratio that enables other elephants (both bulls and cows) to distinguish both the maturity of the bull concerned as well as its phase of musth and respond appropriately.

Other studies have shown that the length of musth increases as bull elephants mature, with the fittest demonstrating a long mid-phase. The Greenwood *et al.* (2005), studies demonstrated that during mid-musth these older males release 'an optimal ratio of frontalin enantiomers'. Estrous females respond positively to such mid-phase release of pheromones, especially during the time of ovulation, and the musth bulls concerned are therefore able to send a precise message to females in estrus, perhaps helping their reproductive success over other less mature bulls. Since female elephants are in estrus only for about 4 days every 4-5 years in the wild, it is important that bulls find them and mate with them. As Kahl & Armstrong point out, musth helps a mature bull in finding and impregnating a receptive female.

The findings of Greenwood *et al.* (2005) and Rasmussen *et al.* (2002) have important relevance in the dealing with crop-raiding wild elephants in Asia where most of them are bulls and quite a number of them are in musth. An understanding of musth and behaviour of musth bulls would help formulate more effective deterrent measures to mitigate the perennial human-elephant conflict in Asia.

While the misfortunes of the African elephant are due to its tusks for which it is being slaughtered in large numbers irrespective of sex, in Asia given that females do not have tusks and not all males carry them either, elephant poaching may be a minor problem and so poaching cannot be the terminal threat it is in Africa. Nevertheless,

elephants in Asia are being killed as they interfere with agriculture. The long-term future of elephants, outside the protected areas in Asia is inextricably linked to the tolerance of man.

The human-elephant conflict in Sri Lanka is real, and it is leading in just one direction: the destruction and eventual elimination of elephants, males in particular, from agricultural areas, unless innovative measures are adopted to address the legitimate concerns of the farmers. The management of human-elephant conflict has to be integrated into a proper land-use policy and also must recognize the elephant as an economic asset to the community. Unless people value living with elephants, the slaughter will go on. If the local people could perceive the elephant as an economic asset instead of as an agricultural pest, they will tolerate it on their land. One way that local people can benefit from the elephant in their midst is from revenues it generates, whether through small-scale ecotourism or from projects that help manufacture paper from dung, produce biogas from dung, or promote organic farming using dung.

The human-elephant conflict has replaced poaching as the biggest threat to the elephant in Asia. While the international conservation organizations are concerned over the trade in ivory, non-tuskers or *makhnas* continue to be slaughtered in large numbers. The debate over elephants is an emotional one, between the preservationists and the pragmatists. The problem with wildlife is that the people who wish to preserve it, are rarely those who have to bear the cost. Given that the human-elephant conflict is already bad today, it may become worse tomorrow. Even if we cannot eliminate the conflict altogether, we need to reduce it to tolerable levels. This is the biggest challenge facing the AsESG.

Charles Santiapillai

References

Greenwood, D.R., Comeskey, D., Hunt, M.B. & Rasmussen, L.E.L. 2005. Chirality in elephant pheromones. *Nature*, **438**: 1097-1098

Kahl, M.P. & Armstrong, B. D. 2002. Visual displays of wild African elephants during musth. *Mammalia*, **66**: 159-171.

Poole, J.H. 1987. Rutting behavior in African elephants: the phenomenon of musth. *Behaviour*, **102**: 283-316.

Rasmussen, L.E.L., Riddle, H.S. & Krishnamurthy, V. 2002. Mellifluous matures to malodorous in musth. *Nature*, **415**: 975-976



Photo: Dr. H. I. E. Katugaha

Genetic assessment of Borneo elephants: origin and conservation implications

Prithiviraj Fernando, John Payne, Geoffrey Davison, Raymond J. Alfred, Michael Stuewe and Don J. Melnick.

Abstract The origin of elephants on Borneo has been controversial, with competing suggestions that they were introduced by humans, or indigenous to the island. Genetic analysis of Borneo elephants and comparison to other Asian elephant populations across their range has shown that they are genetically unique and divergent from all other populations (Fernando *et al.*, 2003). The genetic uniqueness of Borneo elephants suggests that elephants colonized Borneo in the Pleistocene and that they have had an independent evolutionary trajectory since then. Here we discuss the possible factors limiting the distribution of the species in Borneo, and the conservation implications of their newly discovered genetic status.

Introduction

The origin of elephants (*Elephas maximus*) in Borneo is controversial. While some authors have suggested elephants were not indigenous to Borneo but introduced by humans (Shoshani & Eisenberg, 1982), others (Deraniyagala, 1950; 1955) have postulated a natural origin. With a land area of approximately 745,000 km^2, Borneo is the third largest island in the world. Sabah, a state in the federation of Malaysia, occupies the north-eastern tip of Borneo, while contiguous and to the south is the Indonesian province of East Kalimantan (Fig. 1). Elephants are restricted to north-east Borneo (eastern Sabah and the northern part of East Kalimantan) within an area approximately 5% of the island. The north-east, south-west orientation of the longitudinal axis of Borneo, makes this area the farthest part of the island from the closest free ranging elephant populations outside of Borneo, in Sumatra and Peninsular Malaysia (Fig. 1).

Prithiviraj Fernando
* Centre for Conservation and Research
35 Gunasekara Gardens
Nawala Road
Rajagiriya
Sri Lanka

John Payne
Geoffrey Davison
Raymond J. Alfred
World Wide Fund for Nature – Malaysia,
Kotakinabalu,
Sabah
Malaysia

Michael Stuewe
Asian Rhino and Elephant Action Strategy Programme
World Wildlife Fund
Washington DC,
USA

Prithiviraj Fernando
Don J. Melnick
Center for Environmental Research and Conservation
Department of Ecology, Evolution, and Environmental Biology
Columbia University
1200 Amsterdam Avenue
New York, NY 10027
USA

*CCR is a member of the Wildlife Trust Alliance

The earliest detailed description of Borneo by a visitor to the island is that of Antonio Pigafetta, the Italian chronicler of Magellan's Spanish fleet, which sailed into Brunei on the north-west coast of Borneo, in 1521. Pigafetta's group was taken to visit the sultan of Brunei on two tame elephants (Harrisson & Harrisson, 1971). There have been no subsequent reports of elephants in Brunei or western Borneo. The next published record of elephants in Borneo is that of Dalrymple, who reported in 1767 that "the eastern part of Unsang abounds with wild elephants". Tanjung Unsang, presumably the Unsang mentioned by Dalrymple, is near the eastern-most tip of Sabah. In 1811 Hunt reported that wild elephants occurred at Kenibatangan (= Kinabatangan), Unsang and Sabahan, and that ivory was traded from Kinabatangan. The reports of Dalrymple (1767) and Hunt (1811) suggest the presence of a large free ranging elephant population in northeastern Borneo in mid 18th century. Ivory from Borneo is reported to have been imported into China in the middle ages (Laufer, 1925). The massive expeditions of Admiral Cheng Ho, which sailed from southern China to South-east Asia in 1405, may have visited the southern Philippines and possibly Kinabatangan (Harrisson & Harrisson, 1971) but did not leave a written record concerning elephants or ivory.

Were elephants introduced?

There are two factors that suggest elephants may not be indigenous to Borneo. Firstly, it is the common belief, of uncertain but long-standing origin, among people who live or work within the Borneo elephant range (J. Payne pers. obs.). Secondly, it seemingly offers a parsimonious explanation for the limited and unusual distribution of the species in Borneo.

Many authors have commented on the possible

anthropogenic origin of elephants in Borneo. Among them, Silva (1968) mentions "...the popular presumption that they are the progeny of elephants given to the Sulu Sultan by the East India Company in 1750..." but provides no reference. Harrisson & Harrisson (1971) summarize: [among] "...the many and conflicting statements we have heard on the subject at various places in Sabah... [during the 1950s-1960s]... a long time ago the Sultan of Sulu, who had previously received some elephants as a gift, decided to move them from his small island and place them on the Borneo coast. The purpose of this act was that these elephants were his representatives in a territory which he claimed to control..." These animals presumably originated from India, as the operations of the East India Trading Company were mostly in India, and it was a center of elephant commerce. Medway (1977) quotes earlier authors (Müller, 1839-1840; Everett, 1893; Banks, 1931) who made similar but unsubstantiated assumptions. Hunt (1811) reported the presence of elephants on Sulu island (southern Philippines), where he spent six months, and he states that these elephants were imported from Banjarmasin (in Borneo) by Banjar settlers. There is also evidence that a flourishing regional trade in elephants existed, with large numbers caught and exported from Sumatra in and before the 16th century (Marsden, 1811) and from the west coast of Peninsular Malaysia in and before the 18th century (Andaya, 1977).

A 'natural' origin?

In antiquity, Borneo formed part of a single land mass referred to as Sundaland, which also included peninsular Malaysia and the islands of Java, Sumatra and many smaller islands of the western Indo-Malayan archipelago (Holloway & Hall, 1998). Sea level fluctuation during Pleistocene glaciations periodically disconnected and reconnected the Sunda islands through submersion and emergence of low lying parts of the Sunda shelf (Holloway & Hall, 1998; MacKinnon *et al.*, 1996). A savannah corridor which allowed faunal and floral migrations from the mainland to the Sunda islands, is thought to have existed between southeast Asia and Borneo during the highly seasonal climates of the late Tertiary and mid-Pleistocene (MacKinnon *et al.*, 1996). Therefore, it is possible that Borneo was colonized by elephants, through land connections with the mainland and other Sunda islands, and that the current population is descended from those early founders.

There have been a few subfossil or fossil remains in Borneo indicative of a pre-historic elephant population. Ribs and a femur of an animal assumed to be *E. maximus*, now in Lambung Mangkurat Museum, Banjarbaru, were dug up in 1987 from a swamp near Banjarmasin, South Kalimantan. They have not been dated but appear to be relatively recent and not fossilized. A portion of a right upper first molar of *E. maximus*, said to have come from a cave in Belait District, Brunei is believed "more likely to be Pleistocene than Holocene" (Hooijer, 1972). Other fossils found in Borneo and reported to be of "elephants" are either extinct genera (*Stegolophodon* and *Palaeoloxodon* teeth of uncertain origin) or tapir (Cranbrook, 2000; Cranbrook *et al.*, 2000). The paucity of *E. maximus* fossils in Borneo is in contrast to the presence of Holocene (< 11, 000 years) fossils (at Niah, Sarawak and Madai, Sabah) of the Malayan tapir (*Tapirus indicus*) and Javan rhinoceros (*Rhinoceros sondaicus*), both of which are now extinct in Borneo. Holocene *E. maximus* teeth have been found in caves in Sumatra (Medway, 1977) and late Pleistocene and Holocene fossils of *E. maximus* in Java (Van den Bergh *et al.*, 1996), suggesting a late Pleistocene colonization of the Greater Sunda islands by *E. maximus*. However, *E. maximus* fossil and subfossil material is rare over much of its range (Maglio, 1973) and the scarcity of such remains in Borneo may be due to chance (Medway, 1977) or may reflect the paucity of zoo-archaeological studies.

Elephant distribution and numbers in Borneo

The earliest records from which distribution data can be gleaned, date back to the late 19th century (St. John, 1863; Pryer, 1881; Jentink, 1884; Anon, 1886; miscellaneous reports in the British North Borneo Herald during the late 19th to early 20th century). The elephant distribution at that time appears to have been similar to that in 1980 with the exception of the Sandakan Peninsula and extreme south-east Sabah, where plantations excluded elephants during 1955-1980 (Davies & Payne, 1982). Thus, apart from the reduction in species range over the past 20 years due to forest loss in Sabah, the distribution of elephants in Borneo appears to have been rather stable as far back as their distribution can be inferred from historical records.

In 1949, the Conservator of Forests, British North Borneo (H. Keith, quoted in de Silva, 1968) estimated the Sabah elephant population size at 2000 animals. His successor, G. S. Brown commented that Keith considered " the actual number may be only half as much or perhaps twice as many" (Deraniyagala, 1955). Contemporary estimates by foresters and planters ranged from 500 to 5000 (Banks, 1949). Davies & Payne (1982) provided an estimate of between 500 (based on known groups in accessible areas) and 2000 (based on known species distribution and likely average population density). Currently, it is estimated that between 1100 – 1600 wild elephants live in Sabah (Asian Rhino & Elephant Action Strategy (AREAS) project, unpublished data), while a much smaller number occur in the northern part of East Kalimantan, contiguous with the Sabah population.

Possible source populations

Transporting elephants by sailing ship across oceans a few hundred years ago would necessitate their boarding, disembarkation and restraint on board, hence a well trained animal. It would also require the transport of large quantities of food and fresh water. Given the logistic difficulties, geographic proximity makes Sumatra

or Peninsular Malaysia the most likely source of animals introduced to Borneo. However, as commonly believed, if they did originate from animals gifted to the Sultan of Sulu by the East India Company, the source population could also be India. Comparison of the Borneo population to the three putative source populations Sumatra, Peninsular Malaysia and India and in addition, to those of Laos, Cambodia, Thailand, Bangladesh, Bhutan, and Sri Lanka in both mitochondrial and nuclear DNA analysis and additionally Vietnam in the mitochondrial analysis, found it to be unique and not closely related to any of these populations (Fernando *et al.*, 2003). Thus, both mitochondrial and nuclear data suggest elephants are indigenous to Borneo and not introduced by humans (Fernando *et al.*, 2003). In the case of mitochondrial DNA, the evidence supporting a 'natural' origin of Borneo elephants was three fold; the uniqueness of the Borneo haplotype, the degree of divergence from other Asian elephant haplotypes, and congruence with the patterns of distribution and relatedness of other Sunda haplotypes.

Although the mitochondrial genome in general accumulates point mutations much more rapidly than the nuclear genome, and the d-loop of the mitochondrial genome faster than other mitochondrial regions, the rate of evolution of the Asian elephant d-loop is only about 3.5% per million years (Fleisher *et al.*, 2001). The earliest time elephants could have been introduced to Borneo by humans is only a few hundred years ago, which would approximate zero in a time scale relevant to mitochondrial evolution. Thus, if elephants were introduced to Borneo, a haplotype identical to that found in Borneo would be found in the source population. The study by Fernando *et al.*, (2003) sampled almost all extant Asian elephant populations across their range but did not observe the Borneo haplotype anywhere else. Therefore, Borneo elephants could not have originated from any extant population.

Given that the Borneo hapoltype was unique, the arguments for an anthropogenic origin of the Borneo population are, 1) Introduction from a source population that has since become extinct or 2) Fixation in Borneo of a very rare haplotype not represented in the range-wide sample. Although the current Asian elephant range is highly fragmented, the extant populations are spread out over a larger part of the historic range. The representative sample analyzed from extant populations in Fernando *et al.*, (2003) is likely to have captured a high proportion of the mitochondrial diversity of the species. The capture probability of a haplotype is directly proportional to its frequency in the population and the size of the sample collected. Therefore, common, rather than rare haplotypes have a greater probability of being 'captured', be it capture of individuals for introduction a few hundred years ago, or collection of samples for study today. If elephants were introduced to Borneo, the founder population would have been only a few individuals and the haplotype/s represented would likely have been the common haplotypes in the source population. Given the study sample in Fernando *et al.*, (2003), it is unlikely that either an extinct source population in which the Borneo haplotype was common, or the introduction of a rare haplotype not sampled in other populations, provide sufficient explanation for a unique Bornean haplotype.

Since 1980, one of us (J. Payne) has been informed many times by people working in the timber and plantation industries in eastern Sabah that elephants were introduced for logging, and have run wild. This suggests that elephants in these forests are those that were released or escaped the logging camps, and their descendents. Indeed, several Thai or Burmese elephants were brought to British North Borneo (now Sabah) by the Bombay Burmah Trading Company around 1948 for log hauling in Tingkayu, south-eastern Sabah and they were all later returned to Thailand because feeding elephants in the hill dipterocarp forests was too problematic (Q. Phillipps & R. Ibbotson, pers. coms.). Therefore, the elephants in eastern Sabah could not be descended from the logging elephants and the study by Fernando *et. al.*, (2003) also supports this. However, it illustrates how an incident within living memory may lead to propagation of a theory in contradiction of historical evidence.

Distribution of elephants in Borneo

As far as can be ascertained from published records, the historic distribution of elephants in Borneo appears to have approximated the current distribution. There is no clear evidence to suggest that elephants were more widespread in Borneo during pre-historic times. Thus, unless fresh evidence is found in the form of sub-fossil or fossil material confirming their pre-historic occurrence in other parts of Borneo, it has to be assumed that their current distribution reflects the past. This limited distribution could be due to ecological or biogeographic factors.

1) Ecological factors: Elephants appear to be largely limited to habitat below 600 m in Borneo and areas of intensive use are below 300 m. Until the widespread commercial logging in the hill and mountain ranges of Borneo started about 30 years ago, the preferred elephant food plants (monocotyledons and pioneer dicotyledonous plants; Olivier, 1978) were very sparsely distributed in these high, predominantly steep regions, and were mainly found along the larger river valleys in the lowlands. The scarcity of suitable food resources in the higher reaches could have limited dispersal and determined elephant distribution. Sukumar (1989) proposed that an Asian elephant needs 75-100 g of sodium daily in order to avoid a deficit. Davies & Payne (1982) and Payne (1992) suggested that availability of sodium strongly influenced the distribution of large mammals, especially elephants and rhinoceros, in Sabah. In Borneo, natural mineral sources such as salt-rich springs and "mud volcanoes", the latter being grey-coloured mineral-rich mud which is forced from underground by methane gas, appear to be more common in the lowlands and are used today

by elephants. Deficiency of minerals in the montane flora and the apparent lack of alternative mineral sources could have influenced elephant distribution.

2) Biogeographic factors: If elephant distribution in Borneo was shaped by the effects of climatic and geologic history, Pleistocene land connections, and geographic barriers to dispersal, we would expect them to similarly affect other taxa, leading to concordant distribution patterns of multiple taxa. Indeed, peculiarities in the distribution of both flora and fauna support the recognition of northeast Borneo as a unique biogeographic unit (MacKinnon *et al.*, 1996). The north-western lowlands are considered much richer floristically than the north-east or south (Ashton, 1972; MacKinnon *et al.*, 1996), and an area concordant with known elephant distribution in Borneo is recognized as the east coast Sabah floristic sub-province (Wong, 1998). This area lies east of a rough arc from Mount Kinabalu along the eastern flank of the Crocker Range southeast through the Sook, Pendawan and Maliau basins to the Tanjung Redeb area of northeast Kalimantan. A number of mammals including two endemic squirrels (*Petaurillus hosei* and *P. emiliae*), two rats (*Chiropodomys major* and a newly discovered species at Danum Valley, Sabah), a mongoose (*Herpestes hosei*) (MacKinnon *et al.*, 1996) and 24 species of bats (Payne *et al.*, 1985) are confined to this region. This is also reflected in the avifauna, with a number of species replacements in Sabah by semispecies or allo-species (black-headed pitta *Pitta ussheri* versus garnet pitta *P. granatina*, white-fronted falconet *Microhierax latifrons* versus black-thighed falconet *M. fringillarius*), several species absences in Sabah where a form that is montane in the rest of Borneo occupies both the montane and the lowland niche of the "missing" Sabah lowland form (grey-breasted spiderhunter *Arachnothera affinis*, black-fronted leafbird *Chloropsis flavocincta*), species absence in Sabah without any replacement form (Hose's oriole *Oriolus hosei*), and presence in Sabah without any replacement form elsewhere in Borneo (fulvous-chested flycatcher *Rhinomyias olivacea*). The latter species and two woodpeckers, the speckled piculet *Picumnus innominatus* and greater goldenback *Chrysocolaptes lucidus*, are examples of species that are widespread in South and South-east Asia, yet—like the Asian elephant—are confined in Borneo to Sabah in the north-east. These examples are all described in Smythies (1999). Thus, the distribution of elephants could be a consequence of biogeographic factors.

It is possible that resource availability and biogeographic factors, or both led to the limited elephant distribution in Borneo. However, elephants are extreme generalists and are able to survive in a wide range of habitats. Also, in many parts of Asia, elephant range extends well over 1000 m [upto 3200 m in NE India (AREAS unpublished data)]. Therefore, while resource availability may have had some bearing on the distribution of elephants, there does not appear to be a clear correlation, and data on resource availability is insufficient to make a compelling argument. In the same manner, the referred to biogeographic patterns are based on species distribution, hence afford only very coarse comparisons. Future genetic analysis of other taxa both widespread and limited in distribution to the northeast, and correlation with their ecological affinities and species history on Borneo could better evaluate the relative importance of the two hypotheses. We also encourage detailed morphometric analysis of Borneo elephants, as well as search for sub-fossil and fossil *E. maximus* material in Borneo to further establish the evolutionary history of Asian elephants in Borneo.

Conservation implications

Deraniyagala (1950; 1955) considered the Borneo elephant a distinct subspecies, based on its appearance, the examination of a type skull in the British Museum and a sub-fossil tooth, originally reported to be of an elephant, from a cave in Bau, upper Sarawak, now in the Sarawak Museum. He considered Bornean elephants to be smaller in stature, have straighter more slender tusks and their skull to be shallower dorsally above the rostrum, than Indo-Ceylon (Ceylon = Sri Lanka) elephants (Deraniyagala, 1950; 1955). Subsequent workers have considered them not to be a distinct taxon, but the same as the Indian elephant *E. maximus indicus* (Shoshani & Eisenberg, 1982) or Sumatran elephant *E. maximus sumatrensis* (Medway, 1977), based on the assumption that they originated from stock introduced by humans or that the morphological differences were insufficient to warrant such a distinction.

The difficulty of delineating subspecies has led to the development of subspecific designations that are more definable and relevant to conservation, such as Evolutionarily Significant Units (ESU) and Management Units (MU) (Ryder, 1986; Moritz, 1994), although they too are not without problems (Paetkau, 1999). In the case of the Asian elephant, based on mitochondrial DNA analysis, the Sumatran population has been suggested to represent an ESU (Fleischer *et al.*, 2001). If Borneo elephants were feral descendants of introduced animals, in terms of Asian elephant conservation they would be of lesser importance than as a naturally occurring unique population at the edge of the species range. Conversely, if they were indigenous, they could represent an ESU or MU depending on their level of divergence, and they would need to be managed separately from other Asian elephants especially in *ex-situ* conservation.

Borneo elephants were monophyletic in the mitochondrial phylogenetic analysis and displayed uniqueness in the microsatellite analysis of Fernando *et al.*, (2003). The independent evolutionary trajectory of Borneo elephants from that of other Asian elephants, fulfill the criteria for recognition as a separate ESU (Fernando *et al.*, 2003). Thus, there is good reason to devote continued efforts to maintain wild breeding populations of the Borneo elephant and their *in-situ* conservation should be a priority in Asian elephant conservation plans. In addition, Borneo elephants should

be managed separately in *ex-situ* conservation with the objective of preserving their unique genetic make up and should not be cross bred with animals originating from other Asian elephant populations. Examination of pedigrees of captive bred animals and planned future mating of selected individuals, as well as avoiding further breeding of animals with mixed parentage, would be desirable.

The extent of genetic variability within a population determines its evolutionary potential and its ability to cope with environmental change, disease, and demographic perturbations. Loss of genetic variability can result in lowered individual fitness, impaired adaptability (Frankel & Soule, 1981; Allendorf & Leary, 1986), and inbreeding depression (Ralles *et al.*, 1986; Charlesworth & Charlesworth, 1987). However, some natural populations appear to thrive in spite of greatly reduced genetic variability (Ellegren *et al.*, 1993) and populations that undergo inbreeding for extended periods could purge deleterious recessive genes from their genomes and reduce so-called genetic load (Lande, 1988). Although outbreeding generally has beneficial effects, outbreeding depression can occur through loss of local adaptation or the dissolution of co-adapted gene complexes (Templeton, 1986).

Borneo elephants display a significantly low level of genetic variability (Fernando *et al.* 2003). A lesser degree of genetic diversity loss in African Addo elephants led to recommendations for introduction of elephants from other populations (Whitehouse & Harley, 2001). However, the loss of diversity in Addo elephants was very recent and human induced. Given the evolutionary history of Borneo elephants, we would advocate caution before any such intervention is considered. Studies of overt indicators of inbreeding depression such as congenital deformities, behavioral changes, lowered fertility, sperm deformities and lowered recruitment would be logical next steps, in the conservation and management of this population. A wider study of the genetic variability of the Borneo elephant, with an extensive geographic sample and analysis of additional nuclear markers such as Major Histocompatibility Complex (MHC) loci, which are directly involved with the immune response, would provide greater insights into their genetic viability. Additionally, studies on the ecology, behavior and social organization of Borneo elephants could provide insights into the adaptation and survival of insular populations of large mammals.




Fig. 1 Map of NE Borneo showing past and present Elephant ranges.

References

Allendorf, F.W. & Leary, R.F. (1986) Heterozygosity and fitness in natural populations of animals. In *Conservation Biology. The Science of Scarcity and Diversity* (ed. M.E. Soule), pp. 57-76. Sinauer, Sunderland MA.

Anon (1886) *Colonial and Indian Exhibition, 1886. Handbook of British North Borneo*. W. Clowes, London.

Andaya, B. (1979) *Perak, the Abode of Grace*. Oxford University Press, Kuala Lumpur.

Ashton, P.S. (1972) The Quaternary geomorphological history of western Malesia and lowland forest phytogeography. In *The Quaternary Era in Malesia* (eds. P. Ashton & M. Ashton), pp. 35–49. Dept. of Geography, University of Hull, Misc. Ser. No. 13.

Banks, E. (1931) A popular account of the mammals of Borneo. *J. Malay. Brch. R. Asiatic Soc.*, **9**(2): 1-139.

Banks, E. (1949) *Bornean Mammals*. Kuching Press.

Charlesworth, D. & Charlesworth, B. (1987) Inbreeding depression and its evolutionary consequences. *Annual Review of Ecology and Systematics*, **18:** 237-268.

Cranbrook, Earl of (2000) Northern Borneo environments of the past 40,000 years: archaeozoological evidence. *Sarawak Museum Journal*, **55** (76, new series): 61-109.

Cranbrook, Earl of, Currant, A.P. & Davison, G.W. (2000) Quaternary mammal fossils from Borneo: *Stegodon* and *Hippopotamus*. *Sarawak Museum Journal*, **55** (76, new series): 215-233.

Dalrymple, A. (1767) *A Plan for Extending the Commerce of this Kingdom and of the East India Company*.

Davies, G. & Payne, J. (1982) *A Faunal Survey of Sabah*. WWF Malaysia, Kuala Lumpur.

Deraniyagala, P.E.P. (1950) The elephant in Asia. *Proc. 5th ann. Sess. Ceylon Assoc. Sci.*, (1949) **3:** 1-18.

Deraniyagala, P.E.P. (1955) *Some Extinct Elephants, Their Relatives and the Two Living Species*. Government Press, Ceylon.

Ellegren, H., Hartmen, G., Johansson, M., & Andersson, L. (1993) Major histocompatibility complex monomorphism and low levels of DNA fingerprinting variability in a reintroduced and rapidly expanding population of beavers. *Proceedings of the National Academy of Science of the USA*, **90:** 8150-8153.

Everett, A.H. (1893) A nominal list of the mammals inhabiting the Borneo group of islands. *Proceedings of the Zoological Society, London*, **1893:** 492-496.

Fernando, P., Vidya, T.N.C., Payne, J., Stuewe, M., Davison, G., Alfred, R.J., Andau, P., Bosi, E., Killbourn, A., & Melnick, D.J. (2003) DNA analysis indicates that Asian elephants are native to Borneo and are therefore a high priority for conservation. *PLoS Biology*, **1:** 1-6.

Fleisher, R.C., Perry, E.A., Muralidharan, K., Stevens, E.E., & Wemmer, C.M. (2001) Phylogeography of the Asian elephant (*Elephas maximus*) based on mitochondrial DNA. *Evolution*, **55:** 1882-1892.

Frankel, O.H. & Soule, M.E. (1981) *Conservation and Evolution*. Cambridge University Press, Cambridge.

Harrisson, T. & Harrisson, B. (1971) The Prehistory of Sabah. *Sabah Society Journal Monograph*, **4, I-xxii:** 1-272.

Holloway, J.D. & Hall, R. (1998) SE Asian geology and biogeography: an introduction. In *Biogeography and Geological Evolution of SE Asia* (eds. R. Hall, & J.D. Holloway, pp. 1-23. Backhuys Publishers, Leiden, The Netherlands.

Hooijer, D.A. (1972) Prehistoric evidence for *Elephas maximus* L. in Borneo. *Nature, London*, **239:** 228.

Hunt, J. (1811) *Some Particulars Relating to Sulo, in the Archipelago of Felicia*. Source not stated, photocopy in the Tun Haji Mohd. Fuad Stephens Research Library, Kota Kinabalu, Sabah, Malaysia.

Jentink, F.A. (1884) De oliphant en de rhinoceros of Borneo. *Tijdschr. Ned. Aardrijks. Genootsch.*, (2)**1:** 554-559.

Lande, R. (1988) Genetics and demography in biological conservation. *Science*, **241:** 1455-1460.

Laufer, B. (1925) *Ivory in China*. Field Museum of Natural History Chicago. Anthropology Leaflet No. 21.

MacKinnon, K., Hatta, G., Halim, H., & Mangalik, A. (1996) *The Ecology of Kalimantan*. Periplus Editions Ltd., Hong Kong.

Maglio, V.J. (1973) Origin and evolution of the Elephantidae. *Transactions of the American Philosophical Society Philadelphia. New Series*, **63**: 1-149.

Marsden, W. (1811) *The History of Sumatra*. Republished in 1966 by Oxford University Press, Kuala Lumpur.

Medway, Lord (1977) *Mammals of Borneo*. Monographs of the Malaysian Branch, Royal Asiatic Society No. 7, Kuala Lumpur.

Moritz, C. (1994) Defining 'evolutionary significant units' for conservation. *Trends in Ecology and Evolution*, **9:** 373-375.

Müller, S. (1839-40) De zoogdieren van den Indischen

Archipel. In *Verhandelingen over denatuurlijke geschiedenis der Nederlandsche overseesche bezittingen Zoologie (Mammalia)* (ed. C.J. Temminck), pp. 1-57. Leiden.

Olivier, R.C.D. (1978) On the ecology of the Asian elephant. Ph.D. thesis, University of Cambridge, UK.

Paetkau, D. (1999) Using genetics to identify intraspecific conservation units: a critique of current methods. *Conservation Biology*, **13:** 1507-1509.

Payne, J. (1992) Rarity and extinctions of large mammals in Malaysian rainforests. In *Proceedings of the International Conference on Conservation of Tropical Biodiversity, "In Harmony with Nature", 12-16 June 1990* (eds. S.K. Yap & S.W. Lee), pp. 310-320. Malayan Nature Society, Kuala Lumpur.

Payne, J., Francis, C., & Phillipps, K. (1985) *A Field Guide to the Mammals of Borneo*. The Sabah Society & WWF Malaysia.

Pryer, W.B. (1881) Animal life in Borneo. *Zoologist*, **5 (3):** 393-398.

Ralls, K., Harvey, P.A., & Lyles, A.M. (1986) Inbreeding in natural populations of birds and mammals. In *Conservation Biology. The Science of Scarcity and Diversity* (ed. M.E. Soule), pp. 35-56. Sinauer, Sunderland MA.

Ryder, O.A. (1986) Species conservation and systematics: the dilemma of subspecies. *Trends in Ecology and Evolution*, **1:** 9-10.

Shoshani, J. & Eisenberg, J.F. (1982) *Elephas maximus. Mammalian Species*, **182**, 1-8.

de Silva, G.S. (1968) Elephants of Sabah. *Sabah Soc. J.*, **3:** 169-181.

Smythies, B.E. (1999) *The Birds of Borneo*. 4th edition, revised by G.W.H. Davison. Natural History Publications, Kota Kinabalu.

St. John, S. (1863) *Life in the Forests of the Far East. Second Edition.* Smith, Elder, London.

Sukumar, R. (1989) *The Asian Elephant. Ecology and Management*. Cambridge University Press, Cambridge.

Templeton, A.R. (1986) Coadaptation and outbreeding depression. In *Conservation Biology. The Science of Scarcity and Diversity* (ed. M.E. Soule), pp. 105-116. Sinauer, Sunderland MA.

Van Den Bergh, G.D., Sondarr, P.Y., De Vos J., & Aziz, F. (1996) The proboscideans of the South-East Asian islands. In *The Proboscidea: Evolution and Palaeoecology of Elephants and Their Relatives* (eds. J. Shoshani & P. Tassy), pp. 240-248. Oxford University Press, Oxford, UK.

Whitehouse, A.M. & Harley, E.H. (2001) Post-bottleneck genetic diversity of elephant populations in south Africa, revealed using microsatellite analysis. *Molecular Ecology*, **10:** 2139-2149.

Wong, K.M. (1998) Patterns of plant endemism and rarity in Borneo and the Malay Peninsula. In *Rare, Threatened and Endangered Floras of Asia and the Pacific Rim* (eds. C.I. Peng & P.P. Lowry), pp. 139–169.



Photo: Charles Santiapillai




Photo: Heidi Riddle for International Elephant Foundation (IEF)

Establishing a sustainable model for the long-term conservation of the elephant in Sri Lanka

Ravi Corea

The conservation of large vertebrates will be one of the biggest challenges that would be faced by conservationists in the new millennium. Shrinking habitats mainly due to human population pressures are creating a situation that is making even the most ardent conservationists feel current efforts can end as dismal failures. A reoccurring lament is how could the Asian elephant (*Elephas maximus*) be saved in numbers large enough to sustain it amidst such a discouraging and impossible situation?

The last large populations of the endangered Asian elephant share space with some of the most marginalized groups of people in the elephant's range countries. For example, in Sri Lanka the population is 78 percent rural and the elephant shares space mostly with rural farmers. Marginal lands are increasingly brought into agriculture production even though agriculture is the least dynamic sector of the economy, accounting for only 19.4% of GDP in 2001 as shown by a survey conducted by the Asian Development Bank. So nearly 70% of rural farmers are marginalized and live in significant poverty. A Participatory Rural Appraisal conducted by Sri Lanka Wildlife Conservation Society (SLWCS) sociologist, Zeenath Khalid of the Weheragalagama village in the Wasgamuwa region showed that 65% of the population moved in a constant debt cycle. Such low and unpredictable incomes are key features of poverty.

Adding to the considerable suffering of these people is the human-elephant conflict. Human-elephant conflict is perhaps one of the biggest environmental issues Sri Lanka is facing today. Elephants being the largest terrestrial mammals frequently range outside the borders of even the largest national parks in Sri Lanka. Thus setting aside enough habitats to support a large population of such a highly mobile animal as the elephant is almost impossible. Current research conducted by the SLWCS in the Wasgamuwa region shows that there are more elephants outside the Wasgamuwa National Park substantiating the claim that nearly 70% of the elephants in Sri Lanka roam outside the protected areas. Obviously the number of elephants that can be supported by a conservation area and its buffer zone will depend on the tolerance of the people who share their land with the elephants. People will tolerate elephants in their backyard only if they can reap tangible benefits from their presence or if they are provided protection from the frequent elephant attacks on their crops, property and lives or adequately compensated for their losses. Ensuring the long-term survival of such a large and highly mobile animal as the elephant outside the network of protected areas though difficult is critical for the long-term conservation of the elephant. It is also highly unlikely that one government department will ever have the capacity to handle an issue of such magnitude alone. Private-public partnerships will play an important role in ensuring a future for the Sri Lankan elephant in the new millennium. At the government level, there must a be sincere effort to appreciate and accommodate the contributions of private organizations and individuals for elephant conservation. Policies must be developed to encourage such public-private partnerships. Collaboration and sharing of information between individual researchers and conservation organizations is also vital to support these efforts. Additionally the international donor agencies that support elephant conservation should provide incentives for such collaborative efforts and public-private partnerships.

There is a need to develop an innovative approach to elephant conservation in Sri Lanka if elephants are to survive in significant numbers outside the system of protected areas. At present there are no incentives at all for rural people to support elephant conservation.

In the efforts to conserve the elephant some of the biggest and pressing issues are:

1. How can the need to conserve and protect the elephant be reconciled with issues such as human elephant conflict and with the needs and aspirations of stakeholders who are most impacted by elephants.

2. How to develop economically and logistically feasible solutions that are sustainable as well as support the lifestyles and culture of the people of an area.

3. How to protect the last remaining habitat of elephants outside the protected areas.

At present most conservation measures have been developed as reactive measures in response to the proximate issues of elephant conservation, which are: protection, management and establishment of protected areas. Most of these efforts though have been applied in an *ad hoc* manner and have not contributed greatly to conserve the elephant, or address the ongoing issues of human elephant conflict. There is also an urgent need to apply conservation measures in a proactive manner especially in areas where conflict is still low in intensity

Ravi Corea
Sri Lanka Wildlife Conservation Society
127, Kings Land Street,
Nutley NJ 07110, USA.

and there is still opportunity to set benchmarks for elephant conservation. An example of such an approach to elephant conservation is the Somawathiya Chaitiya Temple Project of the SLWCS, which was initiated this year.

Another mostly ignored issue is how to sustain projects that are initiated with donor funding. There is a vital need to understand that while donor funding is critical to keep the processes of conservation moving forward and to initiate new projects, it would be a misjudgment to believe that this periodic infusion of donor funds dispersed annually would suffice to protect elephants or help maintain projects over the long term. Currently there is a tendency to focus solely on projects and their immediate outcomes and ignore other issues such as:

Institutional infrastructure
Individual researcher capacity
Operations and administrative overheads
Institution/individual capacity building
Long-term funding
Sustainability

As much as there is a need to help the elephant, there is a critical need to help the individuals and organizations that are committed to long-term *in-situ* elephant conservation.

A good example is the efforts to mitigate human elephant conflict. There is no definite pathway to resolve human-elephant conflict one hundred percent even with community participation. Considering it is a human endeavor, these efforts are always challenged by new concerns, issues and problems. The whole human-elephant conflict resolution effort is an exercise in incremental learning where we are constantly learning new things about working with rural communities as well as about the ingenuity of elephants. So it is necessary to have an adaptive management approach to apply lessons as they are learned through ongoing research efforts. The most important lesson the SLWCS has learned is that you cannot just erect an electric fence or initiate a project and then walk away. It is important to be involved in the project throughout either directly or indirectly so as to make sure there is accountability, as well as monitoring and evaluation of these efforts continuously over time, as well as to make sure that these efforts stay current by applying new concepts as they are developed from lessons learned in the field. For an individual researcher or an organization to invest such an amount of time on a project means they need to have the resources to keep them in operation over the long term. This lack of operational support is what has curtailed the success rate of efforts to conserve the elephant in the wild as well as to resolve human elephant conflict successfully.

While there is heightened awareness internationally in regard to the need to conserve both the African and Asian elephant – there seems to be a conflict of interest between *in-situ* and *ex-situ* conservationists as to how to move forward in this regard. International institutions such as zoos, safari parks and circuses have been investing most of their resources on *ex-situ* conservation efforts, obviously in an attempt to circumvent the increasing global objection to the capture of wild elephants for trade, they have a critical need to develop a sustainable captive population to fulfill their needs. The irony of this is that for conservationists who are working on *in-situ* elephant conservation, breeding elephants is not an issue – in fact elephants breed too well in the wild contributing to the overall problem. The issue for *in-situ* conservationists is to save as much of the remaining habitat available for elephants especially outside the protected areas. So while zoos, safari parks and circuses are investing millions of dollars on developing artificial insemination techniques and trying to understand the estrus cycle of female elephants, the *in-situ* conservationists receive only a fraction of this amount for their efforts to save the last wild elephants and their habitat. A major effort must be made to reconcile and integrate the efforts of both parties so that the captive elephants in these institutions can be a conduit to channeling more resources for the *in-situ* conservation of their wild cousins. Other international donors such as WWF, CI, etc., are more interested in safeguarding their corporate images than actually funding *in-situ* conservation initiatives! False labeling has become pandemic - where stickers with the corporate emblems of these organizations are widespread even in countries where they are not active giving a false impression that elephant conservation is receiving more support than it is really getting.

The Sri Lanka Wildlife Conservation Society (SLWCS) for the past 8 years have been developing a sustainable *in-situ* conservation model for elephant conservation in Sri Lanka under its flagship project, Saving Elephants by Helping People (SEHP). The SLWCS practices a holistic approach to conservation. The Society helps local people who are the most affected by wildlife and who in turn most affect wildlife since if we are to effectively conserve nature through research based conservation strategies, then we need to create local support for these projects and programs. Even after eight years the SEHP project stands out as one of the most successful attempts to resolve human elephant conflict in Sri Lanka in an area where humans and elephants share space. By integrating economic incentives such as nature, recreational and responsible tourism to its projects SLWCS is generating revenue to attain sustainability to its elephant conservation, research and community development efforts. The revenue generated helps to:

- Maintain field projects
- Employ local field staff

*The advantages of training villagers as opposed to employing non-locals as field assistants are:

- Facilitation of long-term monitoring and surveying of wildlife by providing resident researchers.
- Capacity building increases the ability of villagers to gain revenue through conservation.
- Increased environmental awareness and appreciation

of nature by locals who are otherwise likely to have adverse impacts on the environment.
- To obtain local support for the entire project and avoid conflict that often arises between local habitants and non-local scientists as a result of their differing interests.
- Reduced costs in conducting long term field research
- Maintain electric fences
- Defray costs of community development programs
- Develop education and awareness programs
- Provide micro loans
- Set up community cooperatives for sustainable development
- Establish insurance and compensation schemes
- Develop grassroots environmental organizations.

The SEHP project integrates ecological research, applied conservation, community participation, community development and sustainable economic development. It is the first community based elephant conservation and research project in Sri Lanka and has received continuous support from the Asian Elephant Conservation Fund of the US Fish & Wildlife Service, Disney Wildlife Conservation Fund, International Elephant Foundation and from Sanjay Kumar and Computer Associates, Inc., of the USA. While the SEHP project is the longest operating program of the Society, the SLWCS has several additional projects for elephant conservation that are at various stages of operations and development.

A project to cultivate crops such as citrus that are not desired by elephants but still have the potential to bring farmers a decent income is in the developmental stages. By encouraging the cultivation of such crops it is possible to buffer farmers from total economic ruin when their fields and homes are raided and damaged by elephants.

The SLWCS has just completed Phase 1 of a broad ranging program at Somawathie Chaitiya National Park with funding support from the Alexander Abraham Foundation. Phase 1 was building an electric fence to avoid the eventual conflict that would have resulted between the increasing number of pilgrims and elephants. The Somawathie Chaitiya temple is 2300 years old and is one of Sri Lanka's oldest temples, lying just north of the Polonnaruwa Archeological/UNESCO World Heritage site and is situated right in the middle of a national park, which consist of prime elephant habitat. The SLWCS built an elephant friendly fence, which does not prevent the elephants from moving in and around the temple premises and Chaitiya but prevents them from harming pilgrims and temple property. The Somawathiya Project offers an incredible opportunity to apply proactively all the knowledge, experience and skills SLWCS has gained from its ongoing elephant research and human elephant conflict resolution projects.

At the request of the Department of Wildlife Conservation a project was initiated recently to resolve human-elephant conflict at one of Sri Lanka's smallest national parks, the Lahugala Kitulana National Park, which is 1,500 ha in extent but has one of the largest populations of elephants numbering over 250. The US Fish & Wildlife Service, Disney Wildlife Conservation Fund and the International Elephant Foundation are providing the funds for this project. Lahugala will be the culminating point of all the experiences that the SLWCS has gained from its community-based human elephant conflict resolution projects. The Society is bringing together for the first time in Sri Lanka the national and local government officers such as the Divisional Secretary, Department of Wildlife Conservation, Grama Niladharis, · Police Department; Provincial Administrators, Provincial Council officers & members, villagers and village level societies to develop and maintain a human elephant conflict resolution project.


To achieve environmental conservation goals today it has become important to move away from a purely protectionist method of conservation to a multidisciplinary science. The final outcome of our collective efforts will be measured by how well we have balanced the needs of people while at the same time preserving nature. At the same time it is imperative that conservation must be practiced at a scale that will provide protection to the most diversity of species and ecosystems to ensure sustainable livelihoods for people over the long term. The Sri Lankan elephant is a flagship for the diminishing biodiversity of Sri Lanka. It is also an inherent artifact of Sri Lankan culture, folklore and religion thereby the elephant is a representative of the cultural traditions and history of Sri Lanka. By focusing attention on the elephant's plight we are providing protection to a multitude of other species, habitats, ecosystems as well as conserving some of the cultural traditions of Sri Lanka.

For the past one-year the SLWCS has been integrating responsible travel initiatives to make its projects sustainable over the long term as well as to develop economic incentives for the local people at our project sites to support as well as benefit from the elephant conservation and research efforts of the Society. All these efforts fall into the goals SLWCS have for long-term elephant conservation work in Sri Lanka. We are stressing that community development and sustainable economic development must be ultimate goals that coincide with our conservation and scientific research efforts. The education and sustainable economic development of rural communities is imperative for the co-existence of both elephants and humans over the long-term. If wildlife conservation is to really work then it must move from the administration, management and operations domain of private and public conservation institutions into the hearts, minds, outlook and aspirations of local stakeholders. SLWCS projects offer great opportunities for regular people to get involved in actual field research, *in-situ* conservation, community development and capacity building programs. Further information on how to get involved is available in the Society's website at www.SLWCS.org.

Acknowledgements


The Sri Lanka Wildlife Conservation Society is extremely grateful to the following organizations and individuals who have supported us financially and believed in the vision we have for sustainable wildlife conservation in Sri Lanka. It is mainly their support that helped us to be successful in our conservation efforts. We sincerely thank the Disney Wildlife Conservation Fund, ASPCA, Computer Associates International Inc., Asian Elephant Conservation Fund of the US Fish & Wildlife Service, International Elephant Foundation, the Leo Model Foundation, UNESCO, United Nations Foundation, Travellers Conservation Foundation, the Abercrombie & Kent Global Foundation, the Rufford Maurice Laing Foundation, the Born Free Foundation, The Nature Conservancy, Cleveland Metro Park Zoo and the Alexander Abraham Foundation, and individuals such as Sanjay Kumar and Sir Arthur C. Clarke for giving us grants to conduct our projects as well as to develop the Society. Additionally we thank the numerous people who had volunteered their time or had worked for us and made available their knowledge and expertise over the years to make our projects successful. A special thank you to our village hosts, who by participating in our projects ensured their success.

On a personal note, I thank the Center for Environmental Research & Conservation at Columbia University and its Director, Dr. Don Melnick for giving me the opportunity to study and further my knowledge of Conservation Biology. The Wildlife Trust, USA and its Director, Dr. Mary Pearl for giving me the first grant to initiate my elephant conservation efforts, and Dr. Josh Ginsberg of the Wildlife Conservation Society, USA for being supportive of my conservation and education efforts. I especially like to thank the former Member of Parliament and Governor of the Central Province, the late Honorable Monty Gopallawa for his unstinting support to SLWCS and its projects. I warmly thank Lyn de Alwis, Jayantha Jayewardene and Dr. Charles Santiapillai for their advice and continuing support from even before the establishment of the Saving Elephants by Helping People project and Dr. Natarajan Ishwaran for his continuing support with the current conservation efforts. I also like to say a sincere Thank You to all the SLWCS staff members without whose dedication and commitment none of this would have been possible.



Before the electric fence there was no boundary to separate elephants and humans





The fence provides a non-lethal and safe boundary between elephants and humans



Artist: Manjula Wijesundara

Survey of Elephants in the Mannar District, Sri Lanka

S. Wijeyamohan, S.R.B.Dissanayake and Charles Santiapillai

Participants who assisted in the survey of wild elephants in Mannar District: Ms. A.S. Rangini, Ms. A.A. Hirosha, Ms. R.K. Nishanthy, Ms. P. Ahalya, Ms. S. Vasuki, Mr. J.S. Bernard, Mr. T.M. Sajithran, Mr. V. Ravi, Mr. U. Ragulan, Mr. K.V. Vijithan, Mr. S. Kajendran, Mr. T. Chawchangan, Mr. S.L. Abdul Majeed, Mr. T. Sumanenthiran, Mr. V.S.S. Pragash, Mr. G. Jayaseelan and Mr. S. Krishnakumar (from University of Jaffna, Vavuniya Campus). Ms. Nilmini Jayasena, Ms. H.R.N. Jinadasa and Mr. M.U.A.K. Munasinghe (from University of Peradeniya, Faculty of Veterinary Science). Dr. T. Prasad and Dr. V. Perera (from the Department of Wildlife Conservation, Sri Lanka).

Introduction

The Mannar region is bounded by 8° 30' N latitude and 80°30' E longitude (Eriyagama 1961). The Mannar District with a total land area of 1,985 km^2, extends from the Moderagam aru in the south to beyond Pali aru in the north. It was and still is a part of Vanni (Lewis 1894). The Mannar District is bounded on the west by Indian Ocean, to the north by Kilinochchi District, to the south by Puttalam District, and to the southeast by Anuradhapura District, and to the northeast by Mullaitivu District and to the east by Vavuniya District. The Mannar District consists of the following four Administrative Divisions: (i) Mannar, (ii) Mantai, (iii) Musali and (iv) Nanadan (Fig. 1). Of these four Divisions, Mantai is administered by the Liberation Tigers of Tamil Eelam (LTTE). The human population is estimated at 140,000, distributed in 528 villages. With a population density of 71 people per km^2, Mannar District is also one of the sparsely populated areas in Sri Lanka.

Historically Mannar District has been the gateway to Sri Lanka. Mannar itself was known in King Solomon's times. The Arab trade between the East and West of India was transacted through the channel in Mannar. Vijaya and his band of followers who constituted the first Sinhalese, are said to have sailed down the west coast of India from Supparaka (north of present day Bombay) and landed in Sri Lanka at Tambapanni, which according to Dipavamsa (the oldest of the Ceylon chronicles) was 'on the most lovely south bank of the river'. The river in question is the Kadamba nathi (Aruvi aru or Malwattu oya) along which Vijaya and his followers traveled to Upatissagama and finally reached Anuradhapura (Nicholas, 1959).

As Brohier (1960) points out, centuries before Vijaya and his Aryan followers reached Sri Lanka, the coastal region of Mannar was 'a great commercial emporium', which was visited by Phoenicians from the Red Sea. The city of Mantai (or Matottam) on the mainland had an extensive trade link with Rome and Greece in the 2nd century A.D, but the archaeological evidence, according to Weerakkody (2000) goes to show that Mantai served as the regular port for foreign trade and travel from as early as the 5th century B.C to the 12th century A.D. and beyond. Ptolemy's account of Ceylon in the 2nd century A.D. names it as 'Margana'. The main exports from Mantai were the famed pearls, precious stones, cinnamon, spices, shanks and elephants (Nicholas, 1959). Pearl fishery in Sri Lanka was the most ancient industry in the world, carried on for 2,000 to 3,000 years (Vane, 1887). According to Abayakoon (1950), Cleopatra's pearls may have come from Mannar, for the Mannar coast was known in Tamil as Muttukarai (pearl coast) in reference to the rich pearl banks or paars found in the shallow sea from Negombo to Mannar in general, and between Marichchikaddu and Silavaturai in particular. Coins of Claudius and Annus Plocamus were used in the purchase of pearls by Romans (Abayakoon, 1950). Robert Knox, the celebrated English prisoner of the last king of Kandy escaped from captivity and arrived at Arippu on the west coast.

Climate

With more than six arid months, the Mannar District is situated in one of the driest areas in Sri Lanka. During the dry season, which extends from mid-March to mid-October, daytime temperatures of well over 30°C (or 85°F) have been recorded. The hottest month is May, when the southwest monsoon is at its height in the south and central hills. The coldest months are December and January coinciding with the northeast monsoon during which the region receiveds between 760 and 1000 mm (or

S. Wijeyamohan
Department of Biological Sciences,
Faculty of Applied Science,
University of Jaffna,
Vavuniya Campus, Kurumankadu,
Vavuniya, Sri Lanka

S.R.B.Dissanayake
Department of Wildlife Conservation,
18 Gregory's Road,
Colombo 7, Sri Lanka

Charles Santiapillai
Department of Zoology,
University of Peradeniya,
Sri Lanka




Fig. 1. The four Administrative Divisions of the Mannar District.

30-40 inches) of rain from mid-October to mid-February. The wettest month is November. Erratic showers are not uncommon during the southwest monsoon, but such rainfall has hardly any appreciable effect on the vegetation as the prevalent heat and comparatively lower humidity result in rapid evaporation (Eriyagama 1961).

Physiography

Much of the land area is flat, being less than 3.65m (12ft) altitude above sea level. Along the coast are sand dunes but towards the Giant's Tank, the landscape takes on a gently undulating aspect, which is more pronounced in the northern direction (Eriyagama, 1961). A few rivers drain the area, and they are from south to north, (i) Uppu aru (Modergam aru), (ii) Kal aru (or Cheddikulam aru, a river formed of two streams which rise in the high ground of the southern corner of Kilakkumulai South), (iii) Aruvi aru, (iv) Nay aru, which enters the sea to the south of Vidattaltivu, (v) Paranki aru, which reaches the sea at Illupaikadavai and (vi) Pali aru, which flows almost due west, forming the northern border of Mannar District and the southern boundary of Tunukkay division (Fig. 2). It falls into the Gulf of Mannar in the extreme north of Mannar District (Lewis, 1895). Except for Moderagam aru and Aruvi aru, all other rivers become bone dry during the peak of the dry season. A channel from Aruvi aru feeds the Giant's Tank – the largest irrigation reservoir in Mannar District, while another goes to feed the Ahathimurippu tank. Irrigation in Mannar District takes the form of hundreds of man-made reservoirs or tanks scattered all over the country to catch as much water as possible during the short rainy season (Fig. 3). As Lewis (1895) points out, these tanks for the most part form a connected series, one above the other, those lower down receiving the surplus water of the higher tanks. Water from Giant's Tank feeds 160 other lesser tanks before reaching the seas, and only about 4,000 million cubic metres of water finally reach the sea.

Soil

Except for the narrow sandy coastal strip, much of the Mannar District is composed of soils that are grayish-brown sandy clay to sandy clay loams with concretions of carbonates of calcium, sodium, and manganese oxide (Eriyagama, 1961). This fine clay tends to become stony-hard during the drought, making it extremely impervious to water. Given the combination of high impermeability and the high content of sodium, the soils are characterized by a high salt content and high pH (i.e. highly alkaline). Murunkan clay in particular is well known for its excellent qualities to support agriculture. Further inland, soils are reddish-brown in colour. Such soils are well drained and are ideal for agriculture. According to Boake (1888) the subsoil of loam or clay is very fertile, producing most of the timber trees indigenous to the island. Some fields yield 30-fold of paddy and tobacco.

Vegetation

Despite the arid climate and inhospitable landscape, a substantial part of Mannar District is still forested (Fig. 3). The area is rich in biological diversity. Given the arid climate and edaphic conditions, the region has its own peculiar fauna and flora. Along the coastal sand dunes can be seen patches of *Spinifix littoreus* in association with *Ipomoea pescaprae*. Suriya (*Thespesia populnea*) and Malitan (*Salvadora persica*) are two of the commonest trees in the coastal areas of the dry zone. The principal vegetation type is thorn-scrub, which provides some sort of ground cover over much of the region for the soil against excessive loss of water through direct evaporation (Eriyagama, 1961). Many of the plants show marked xerophytic characters such as thorns and spines to protect them against grazers and browsers and also to keep transpiration to a minimum. Much of the island of Mannar is covered with Palmyrah (*Borassus flabellifer*) and Coconut (*Cocos nucifera*) palms. As Eriyagama (1961) points out, given the economic importance of the coconut, it has to a limited extent, replaced palmyrah. The coastal vegetation is characterized by stunted cockspur-thorn (*Acacia eburnea*) and buffalo-thorn. The scrub vegetation consists mainly of species such as *Cassia auriculata, Phoenix zeylanica, Euphorbia antiquorum, E. tirucalli, Calotropsis gigantean*, and *Glycosmis pentaphylla*. The ground vegetation consists of *Aloe vera* var. *littoralis* (Komarika) and *Opuntia spp*. Much of the former extensive mangroves that bordered the causeway between the mainland and Mannar island has been destroyed as a result of almost 20 years of civil conflict. *Rhizophora spp.* with their characteristic stilt roots and numerous pneumatophores are still common in swampy areas, while the ground flora is mostly of the fleshy *Salicornia brachiata* and *Tamarix gallica* (Eriyagama, 1961).

On the mainland, the subsoil of loam or clay being very fertile, most of the economically valuable timber species thrive here. These include Palu (*Manilkara hexandra*), Wira (*Drypetes sepiaria*), Buruta (*Chloroxylon swietenia*), Kohomba (*Azadirachta indica*), and Ebony (*Diospyros ebenum*) typical of the dry evergreen forest. Between Oddaipelankulam and Marikapelankulam on the coastal road to Pooneryn, there are, according to Eriyagama (1961) vast extents of damana land and semi-arid thorn scrub, mainly of cockspur-thorn (*Acacia eburnea*), associated with Murunkan soils. The riparian dominants are the Kumbuk (*Terminalia arjuna*), Kon (*Schleichera oleosa*), and Siyambala (*Tamarindus indica*). The Baobab trees (*Adansonia digitata*) that are found in Mannar and on the mainland are not indigenous to Sri Lanka but were introduced by Arab traders from Africa several centuries ago.

Biological diversity

Mannar District is also rich in mammalian and avian biodiversity. According to IUCN (1990), over 30 species of terrestrial mammals have been recorded but additional




Fig. 2. The principal rivers (i) Modergam aru, (ii) Kal aru, (iii) Aruvi aru, (iv) Nay aru, (v) Paranki aru, and (vi) Pali aru, tanks (dark shading) and sanctuaries (hatched areas) in the Mannar District.

species of Rodents and Bats are present. Threatened animals include the Asian elephant (*Elephas maximus*), sloth bear (*Melursus ursinus*), leopard (*Panthera pardus*), and the water buffalo (*Bubalus bubalis*). A variety of herbivores, such as the spotted deer (*Axis axis*), sambar (*Cervus unicolor*), barking deer (*Muntiacus muntjak*), mouse deer (*Tragulus meminna*), wild boar (*Sus scrofa*), and the black-naped hare (*Lepus negricollis*) occur in the District. In addition, all four species of primates are found here, including the Slender Loris (*Loris tardigradus*). In the coastal waters of Mannar is found one the most seriously endangered large marine mammals, the dugong (*Dugong dugon*).

The villu ecosystem supports a variety of resident and migratory waterfowl, including large breeding populations of painted stork (*Mycteria leucocephala*) and open-billed stork (*Anastomus oscitans*). Other wetland species include garganey (*Anas querquedula*), pintail (*A. acuta*), whistling teal (*Dendrocygna javanica*), spoonbill (*Platalea leucordia*), white ibis (*Ardea melanocephala*), large white egret (*Egretta alba*), cattle egret (*Bubulens ibis*), and purple heron (*Ardea purpurea*). Indian darter (*Anhinga melanogaster*) is abundant in tanks with high fish density. In addition, Wijeyamohan *et al.*, (2002) have recorded a minimum of 33 resident species from the Giant's Tank alone. This tank alone supports 16% of the national total of birds resident in Sri Lanka, and at least 74% of all the waterfowl in the island.

Protected areas

Despite its vast area and sparse human population, Mannar District has only two areas under protection: (i) The Giant's Tank (3,941ha), and (ii) The Madhu Road (26,677ha) Sanctuaries (Fig. 2), but they represent 15.3% of the land area in the District. The Giant's Tank is situated on the coastal plain of northwestern Sri Lanka, about 5km from the sea near Murunkan, 18km southeast of Mannar. It was declared a sanctuary on 24 September 1954 (IUCN, 1990). The Giant's Tank sanctuary is built around the ancient reservoir which has a maximum depth of 3.2m and a pH of 7.5. It is one of the largest and shallowest reservoirs in Sri Lanka known for its high productivity in freshwater fish and a rich avifauna, especially waterfowl. Fish production is estimated at about 200 metric tones per year (IUCN, 1990). The Giant's Tank and its adjoining Sanctuary provide one of the key habitats for the Asian elephant.

The elephant in Mannar District

Historically, the Mannar District has been famous for its rich and diverse wildlife, and was particularly known for its elephants. In 1888, the them Assistant Government Agent of Mannar, Mr. W.J.S. Boake, wrote as follows about the elephant, "*The elephant still finds a home here, and during the rainy season makes his appearance in considerable numbers, being driven out of the denser jungles by the flies and mosquitoes, which infest such localities, into the low scrub and open plains by the sea. They are not very numerous as they used to be, but still may be always found in or near this District. One seldom hears of a tusker now, or of a rogue, but rogues have been present in Vilankulam*" (Boake, 1888).

Elephants have been caught and tamed in Mannar District from very early times. They were exported from the ancient port of Mantai which now lies buried under the sea opposite Mannar island. Claudius Ptolemy (c. 121-150 A.D.) refers to this ancient port as the chief outlet for the export of elephants (Nicholas, 1954a). A particular group of Muslims known as Panikkars were adept at catching wild elephants using a noose. Panikkars were originally from Batticaloa and Badulla. Kraaling was not practiced in the north given the heavy mortality such a method entailed. In 1665 only 20 of the 90 elephants that were captured reached Jaffna alive (Nicholas, 1954b).

In 1880 elephants were very numerous in the Nanadan area that they were regularly captured and sold, mainly to India (Boake, 1888). Even then elephants caused great damage to crops and therefore rewards were paid for their destruction. In 1867, elephants to the value of 2,328 sterling pounds were exported (Boake, 1888).

Rationale for the survey

The last "island-wide" survey of elephants that was carried out in June 1993 (Hendavitharana *et al.*, 1994) did not include the north and east of the island because of the civil conflict then raging in the country. Prior to this survey, Norris (1959) arrived at an estimate of 1500 wild elephants in Sri Lanka, but he did not give any details about the distribution of the elephants. Subsequently, the only comprehensive assessment of elephant range and number was that of the team from Smithsonian Institution between 1967 and 1969 during which McKay (1973) estimated between 1600 and 2200 elephants as a minimum figure for the number of wild elephants in Sri Lanka. Even then, much of the northern portion of Sri Lanka, including the Mannar District and the rest of Vanni was not surveyed in any detail. McKay (1973) estimated the total population in this area to be "probably between 200 and 500 elephants". Therefore, as far as elephants are concerned, the Mannar District has never been surveyed in any detail. Hence the need for an update on the range and abundance of elephants here.

Organizing the survey: *the modus operandi*

The survey of elephants in Mannar District is a part of the overall survey of the elephants in the northwestern region of the island organized by the Department of Wildlife Conservation (DWC). We were given the task of organizing the survey in the Mannar District simultaneously with that in the northwest between 25-27 September 2004.

Surveying elephants in the wild is not an easy exercise. It is even more difficult if the distribution of elephants cuts across areas under different administration, as it is in the




Fig. 3. The extent of forest cover in the Mannar District.

Mannar District. During the preliminary meeting that was held at the Department of Wildlife Conservation in Colombo, we were entrusted with the survey of elephants in the Mannar The Government Agent, Mr. V. Visuvalingam reacted positively to a request made by Mr. D. Karyawasam, the Director-General, Department of Wildlife Conservation (DWC) for assistance in the survey, and promptly convened a meeting of the Assistant Government Agents and Grama Niladharis from the four AGA Divisions (Fig. 1) and briefed them of the nature of the survey and requested that all assistance be given to us. He furthermore, drafted a letter to the Area Commander of Mannar District, Brigadier Sarath Fernando to inform him of the proposed survey. Mr. Visuvalingam also made an appointment with the Brigadier and took us to discuss the logistics of the survey. In this way, we kept the Army Commander fully informed of the dates and places where the survey would take place. We also informed the Inspector of Police, Mr. Navarathna C. Kumara of Murunkan of the planned survey and enlisted the help of his staff. The Government Agent discussed the survey with several NGOs and made sure that there were sufficient number of vehicles and drivers to transport the students to their observation points. His house functioned as an interim office and hostel for the participants during the survey. Prior to the survey, a training workshop was held at the Vavuniya Campus of the University of Jaffna, at Kurumankadu, Vavuniya in order to introduce the survey techniques to the students.

Mantai is under the control of the LTTE, and hence without the explicit permission and support of the LTTE the survey could not have been carried out by anyone. Thus, our first task was to explain the nature of the work to the LTTE and enlist their support for us and the students from the Universities to enter the areas under their control and carry out observations during the day and night. The LTTE gave us their full support for the survey to be carried out in areas under their control.

Survey and census

While it may be possible to count all the elephants in an area of 10 km^2 it is a much more difficult task to count them in an area of 2000 km^2 as for example in the Mannar District. Nicholas (1955) quotes Col. Stevenson-Hamilton, the late Warden of the Kruger National Park in South Africa, who stated categorically that it was impossible to make a census or accurate count of animals including elephants in the bush country of Kruger National Park. What is impracticable in Africa, according to Nicholas (1955) "is altogether impossible here". Their skepticism highlights the difficulty of counting even the largest terrestrial mammal. A large area needs a greater amount of time or a large number of people. This raises the problem of over or under counting due to animal movement during the census, and to different observers recording (or ignoring) the same animal (Rodgers, 1991).

Keeping such constraints in mind, the DWC proposed the adoption of the water-hole count as a basis for determining the range and possibly some assessment of the number of elephants in the northwestern region of Sri Lanka.

In small areas where a distinct dry season limits the availability of water for elephants, it may be possible to get a crude estimate of elephant numbers by counting those that visit all the water holes on a single day (Ramakrishnan *et al.*, 1991). The critical assumption here is that all elephants would visit a water hole once during the day. In the survey of elephants in Mannar District, we adopted this method in order to assess the structure of the elephant population, the abundance being a secondary concern. For management of elephants, it'd be more useful for us to know the percentage of tuskers among the bulls, the proportion of calves in the population, and the sex ratio of adults. We hoped that these could be determined through visual observation of the animals at water holes.

Abundance, as Caughley (1977) argues, can be measured in three ways: (i) as the number of animals in a population, (ii) as the number of animals per unit of area (absolute density), and (iii) as the density of one population relative to that of another (relative density). One of the central issues in wildlife management concerns the estimation of population size. In the absence of information on the size and spatial distribution of a population, it would be almost impossible to investigate size-dependent or density-dependent relationships (Williams *et al.*, 2002). Reliable estimates of elephant populations are essential to address some of the important conservation issues. However, as Caughley (1977) points out, "the majority of ecological problems can be tackled with the help of indices of density, absolute estimates of density being unnecessary luxuries".

It is therefore necessary that methods that are adopted to estimate elephant numbers (N) and density (D) meet at least the minimal standards for scientific rigor. Elephants can be estimated either in a **census** operation where all the individuals of a population are counted, or in a **survey** of a sample of the population. Elephants can be counted either from air, or from the ground on foot or from a vehicle in open areas (Direct observations). An estimate of population density, as Jachman (19696) points out, is obtained simply by dividing the number counted by the size of the area censused, and the density figure obtained in this way can then be applied to surrounding areas with similar characteristics, such as soil types and vegetation. Elephant numbers can be estimated based on signs such as dung-piles, tracks, feeding signs etc (Indirect method). This method is useful in areas where it is almost impossible to observe elephants.

Methodology

The survey of wild elephants was carried on two days (25 and 26 September 2004), and it involved the services of 24 participants. On 25 September, the northern part of the




Fig. 4 Elephant observation points: 1. Moodampitty, 2. Kurai, 3. Parankikulam, 4. Sethuvinayagar, 5. Kovitkulam, 6. Palamadu, 7. Sannar, 8. Periyamadu, 9. Palampitty, 10. Thandikulam, 11. Periyavillu, 12. Sinnavillu, 13. Madu, 14. Pandivirichan, 15. Giant's Tank, 16. Pannaivattuwan, 17. Kunjikulam, 18. Uvayadikulam, 19. Puthukulam, 20. Kalliadikulam, 21. Periyamurippu, 22. Periyakattu, 23. Mullikulam, 24. Karadikulam, 25. Kal aru, 26. Anaivillu, 27. Kakkupadayan, 28. Pandarikulam, 29. Eruvettan, 30. Achchankulam.

Mannar District was covered. This area is largely under the control of the LTTE, while on the second day, the southern part of the District was surveyed. Observations were carried out from 1600 to 2300 hrs and beyond in some areas. A number of water holes or (Fig. 4) were identified from where observations on elephants could be carried out. At every water hole, a group comprising 2-3 students was stationed with either a Grama Niladhari or a local villager. A mobile patrol visited all the observation points and covered other areas which were not manned by observers. Direct observations of elephants were the preferred method of investigation. At every sighting of elephants, their number and composition was noted.

The most direct way to estimate the abundance of an elephant population is to count all individuals in a defined area. As Shoshani *et al.*, (2004) point out, counting elephants was done so that only the minimum number of individuals is reported. Duplication was reduced to a minimum or avoided by keeping a tab on the time and distance between successive observations. Wherever possible, animals were classified into calves, juveniles, subadults, adult females and adult males and tuskers. Therefore interviewing local residents is an extremely important aspect of this survey. However, elephants visit water holes frequented by people, more during the night than in the daytime. If no elephants were sighted, then the number of elephants that visited the area most. recently was recorded from the villagers. In areas where there were clear, fresh footprints, their diameter or circumference was measured in order to estimate the approximate shoulder heights of the elephants that visited the area, using the formula that twice the circumference of the front foot gives the approximate shoulder height of an elephant (Boyle, 1929; Sukumar *et al.*, 1988). In Africa, elephant's should height has been estimated indirectly using the hind footprint length since the latter is known to relate to shoulder height, and so it could be used to estimate age distributions in a population (Western *et al.*, 1983). Dung and footprints or other spoor, such as chewed vegetation and scratching posts, provide indirect data (Shoshani *et al.*, 2004).

Results and discussion

There were three problems that faced the observers. The **first** relates to the unexpected rainfall which was very high on the second day of the survey. In Mannar District, the northeast monsoon rains usually commence by mid-October. Hence the survey was planned in advance of the rains so that elephants could be observed, sexed, classified and counted at the few remaining waterholes that are not bone dry. In Mannar District, almost all the waterholes had become dry and the drought was one of the worst in living memory of the people. In the LTTE controlled areas, only one tank (besides the Giant's Tank) had water and that is at Sannar which, according to local people, never dries up. South of the A14 highway, Mullikulam had little water but Periyakattukulam had become dry by the time the survey had started. However, the sudden and unexpected rain (which fell first on 23 September) shifted the movement of the elephants away from the water holes and dispersed them into the forest and along the rivers. Hence the low counts at the observation points on the second day of the survey.

The **second** problem relates to the ban imposed by military on taking binoculars into LTTE controlled areas. As a result of this constraint, it was not possible to sex and classify animals seen at a distance. It is unfortunate that even *bona fide* researchers are denied an opportunity to use the binoculars at a time when the guerrillas all over the world have access to high-powered night vision equipment, telescopes and Global Positioning System (GPS) for surveillance!

The **third** problem concerns the danger posed by unexploded landmines in some areas which poses a serious threat to the life and limbs of people. Such areas could not be surveyed. In the estimation of elephant numbers, every effort was made to discard numbers that may belong to the same population (e.g. in Giant's Tank, a group of 53 animals was not included as it may belong to the much larger group of 77 animals recorded from here).

Distribution of elephants

Elephants in Mannar District enjoy wide distribution that stretches from the Modergam aru in the south to north beyond Mundrampitty near Pali aru (Fig. 5). They are absent along the coast from north of Achchan kulam to just north of Kalliyadi. Despite this large extent of elephant range, the density of elephants varies from place to place, being more in the areas under the LTTE control, just north of the A14 highway. This is largely due to the fact that there are numerous water holes scattered in this area and also substantial forest cover to the east. Elephants are known to move from the south to the north across the A14 highway. However, the appropriation of land along the A14 highway by the Army and the establishment of camps and bunkers, in some places, disrupts elephant movement. Besides, at least 5 elephants have been electrocuted along the A14 highway between Parayanalankulam in the east and Murunkan in the west, during the past three years, through contact with the low-slung naked electricity wires that the army and police use to tap electricity from the national grid.

Abundance of elephants

The results from the survey are given in Table 1. A total of 220 elephants were recorded from the survey, of which 169 were from the LTTE controlled areas, while 51 animals were recorded south of Aruvi aru. In addition, information provided by villagers (and hunters) indicates that up to 350 animals could be using the Mannar District as their habitat during the drought. We should not ignore the knowledge and experience of the local rural people, who in most instances know much more about elephants than urban pundits who just visit an area for a day or two.




Fig. 5 The dry season range of Elephant in the Mannar District.

As Blanc *et al.*, (2003) point out, even the guesses made by people who know the area in question have to be relied upon. Given the closeness of the two estimates of 169 and 189 from the observers and villagers in the LTTE controlled areas for the number of elephants (Table 1), the estimate given by the villagers should be considered an informed guess. Even this figure must be considered a minimum estimate, given the difficulty of counting elephants in forests. A population of 350 elephants in the Mannar District represents approximately 10% of the national total. This translates into a crude density of 0.18 animal per km^2. However, ecological density for the elephant can be much higher in areas of optimal habitat, or during periods of concentration at the height of the drought (Eisenberg & Lockhart, 1972).

The elephant biomass calculations at the crude density of 0.18 per km^2 in Mannar District, work out to be 326 kg per km^2. At ecological densities of 1.0 to 1.2 animals per km^2, the elephant biomass may reach 1,800 to 2,170 kg per km^2. Density, expressed either as individuals per unit area, or as an index to numbers (e.g. number of faecal droppings per 100 stations), is often a valuable and necessary piece of information for management of elephants (Riney, 1982). If measured during the same season in consecutive years, changes in density-related figures will indicate increases or decreases in the population. Biomass refers to the total weight of animals supported by a unit area. The biomass level that a species population sustains represents a relation between the production of food in the environment, and the ability of animals of the species to transform the food into animal biomass (Owen-Smith, 1988).

For such a large mammal as the elephant that can tolerate a lower quality of diet than smaller herbivores, a high proportion of available herbage becomes acceptable food. Density and biomass supply helpful information on what exists at a particular time (present status, standing crop, occurrence).

It appears that there are far more elephants today in Mannar District than was expected. This is not surprising given the resources available to the elephants in Mannar District. In the first place, there is extensive forest still intact in the north. Sri Lanka's forest cover has declined to less than 22% and much of what is left can be found in the north and east. Thus elephants have sufficient habitat, low human population density, and hundreds of man-made reservoirs, a number of which hold permanent water even during the peak of the dry season. The Giant's Tank provides the best area for elephants and other wildlife, especially birds. The LTTE has imposed a strict ban on illegal timber harvesting within the forests in the north. No tree can be cut without their permission. Thus elephants have sufficient habitat to move about and adjust their densities to changing patterns in vegetation.

Population structure

The unexpected rain was in a way, a blessing in disguise in that in some areas, it was possible to see clear footprints which were measured to estimate the approximate shoulder heights of the elephants. A total of 21 elephant footprints in two separate areas were measured. On the basis of the footprint measurements (see Table 2), it appears that the population structure consists of adults (47.6%), subadults (23.8%), juveniles (14.3%) and calves (14.3%). The size classes were estimated based on the criteria given by Eisenberg & Lockhart (1972) and McKay (1973). It is interesting if the observed percentages of the size classes in Mannar District are compared to what was found by Eisenberg & Lockhart (1972) in Wilpattu National Park, where adults comprised 49%, subadults 25%, juveniles 20% and calves 7%. Thus the observations in Mannar District appear to be in keeping with the more extensive censuses performed by Eisenberg & Lockhart (1972). More than 10% of the animals are calves, which is a healthy indicator of recruitment in the population. As Moss (1988) points out, age structures of elephants usually contain peaks and troughs – i.e. there will be numerous individuals in one age class, very few in the next and so on. There is no even distribution of ages through the population. In respect to sex-classes, the male-female ratio is equal until the adult class is reached, whereupon there are only 8% males compared to 31% females (Kurt, 1969). On a separate day, following rains in the LTTE controlled area of Moondrampitty south of Pali aru, we found one very large footprint 39cm in diameter, which is likely to belong to a bull that is about 2.45m in height at the shoulder.

Tuskers

The survey reveals the presence of at least 4 tuskers in the District (Table 1). The proportion of bulls bearing tusks varies enormously among different Asian elephant populations (Sukumar 1989). In Sri Lanka, Deraniyagala (1955) found that 324 bulls out of a sample of 364 were without tusks. i.e. 89% were tusk less or makhnas. McKay (1973) in a much smaller sample of 25 captured bull elephants found 23 (or 92%) were tusk less. In comparison, almost 90% of the bulls are tuskers in South India. It is difficult to determine the % of tusker in the Mannar District based on the results of this survey as it was not possible to determine the adult sex ratio of the animals recorded. However, if we assume an adult male: female sex ratio of 1:3, then the proportion of tuskers in Mannar District would be about 7.3% - exactly the same as that arrived in 1993 census (Hendavitharana *et al.*, 1994).

Human-elephant conflict

The survey also revealed skulls and bones of two elephants but the cause of death could not be ascertained. While there had been much discussion in the media about the impact of wild elephants in agricultural areas in the south, no one highlighted the plight of the people *vis-à-vis* elephants in the LTTE controlled areas.

A key element of elephant conservation is monitoring

Table 1. The number of elephants recorded during the survey. Numbers within [brackets] refer to the approximate number of elephants recorded by villagers just prior to the survey to survey.

location	time	tuskers	adults	subadults	juveniles	calves	total
Day 1 (25 September 2004)							
Parankikulam	1840						2
Parankikulam	1910						13
Palampiddy	1950						[120]
Palampiddy (feeding sign)	1930					1	
Palampiddy (f-prints)	2110						2
Palampiddy (f-prints)	2110						8
Koorai		1	13	7	3	2	26
Giant'sTank	2335	1	43	8	25 (plus calves)		77
Giant's Tank	2350	1	1	1	1	4	
Periyavillu							[30]
Periyavillu (heard)	2025						1
Pandivirichchan(heard)	2340		1				1
Aryvi aru							5
Potkerni							1
Maruthmadhu jct							3
Veppankulam							1
Pitchaivaniparkulam							3
Vilatikulam							[35]
Periyamadukulam							2
Periyamadu							[4]
Periyakattu	2320						15
Total						169 and [189]	
Day 2 (26 September 2004)							
Adaipankulam	1900	1					
Aruvi aru bridge	1715	1					1
Pannivattuwan	2400						1
Pannaivattuwan	0615+			4			4
Pannaivattuwan							[100]
Periyamuripukulam	0600+						1
Thekka pannai		1					[12]
Aathikulam							3
Kallikulam							[19]
Arippu							[30]
Periyakattu	1900						7
Karadikuli	2300						2
Mullikulam							5
Aruvi aru	2220						12
Kal aru (f-prints)	0930						11
Madhu Rd-Mannar A14	2140						3
Total						51 and [161]	
Grand Total	4 tuskers					220 and [350]	

how different forms of human activity influence elephant distribution and abundance (Walsh *et al.* 2001). One of the serious issues that need to be mitigated in the Mannar District is the growing threat of elephants to people and cultivation. The problem has been compounded by the fact that for much of the past two decades, the area was plunged into an internecine civil war and people fled the area to the safety of India as refugees, abandoning their homes, goods and chattel. Today, with the fragile peace still holding between the LTTE and the Government of Sri Lanka, refugees are slowly returning to their original homes to face an unexpected threat – this time from wild elephants. These animals moved into the abandoned home gardens, attracted by the abundance of palatable and highly nutritious food plants. With plenty of food and water, the animals have become 'permanent residents'. Therefore returning refugees have to compete with elephants in cultivating their own land. Elephants killed at least two people and a few were injured but to date no one has received any compensation. The DWC sent some thunder flashes to Mannar District for the first time, so that they could be used to ward off elephants. Elephants can be saved if even a small portion of the money now being used in workshops, symposia, meetings and conferences is directed towards alleviating the hardship of the poor people who bear the brunt of elephant depredations. Conservation has to sustain not only the spirit but also the stomach.

Impact of the civil war on environment

The civil war has radically altered economic, political, and social conditions of the people in the Mannar District with profound impacts on the environment, natural resources and biodiversity. The impact of armed conflict on the environment in Mannar District is perceptible and it occurs through habitat destruction and loss of wildlife, over-exploitation and degradation of natural resources (Shambaugh *et al.*, 2001). Many of the refugees rely on bushmeat for protein. The LTTE has banned the hunting of wildlife except the pig and hare in areas under their control. Outside however, illegal hunting for bushmeat is rampant. At least 50 hunters are involved in the Mullikulam area alone in bushmeat harvesting. Yet bushmeat hunting should not be banned since it would deny the poor people living in remote areas an important source of protein (Whitfield, 2003). It should be regulated. If such harvesting is not regulated, there is a risk that resource depletion and environmental degradation can drag the people into a vicious circle: poverty, further political instability, more armed conflict, greater environmental degradation, and even greater poverty (Shambaugh *et al.*, 2001). Depletion of biodiversity and the natural resource base can undermine the sustainable livelihoods for the people (Santiapillai & Wijeyamohan, 2003).

Conclusion

The survey of wild elephants in the Mannar District represents the first attempt to obtain reasonably reliable minimum estimates of their number. It is a part of a much larger survey carried out simultaneously by the Department of Wildlife Conservation in the northwestern region of Sri Lanka. Until this survey was carried out, there had been no reliable information on the elephant's range and numbers from the north in general and the Mannar District in particular. The status of the elephant varies widely across its range in the island, and moreover its long-term survival is more threatened in some areas than in others. Therefore regional assessment of elephant range and numbers is crucial given that elephants move long distances across political boundaries and protected areas. This is an important consideration in the north since the Vanni is under the control of the LTTE, while all the national parks are under the control of the Government. Decisions made in one area may therefore affect elephant populations in the other.

The water hole count method adopted during the survey is not perfect but useful in areas where the objective is not simply to estimate the number of elephants but more importantly, to understand their distribution, structure and composition. Total counts do not have the advantage of sample statistics to give measures of precision of the estimate (Rodgers, 1991). Besides, the longer the time period taken to do the count, the greater the chances of double counting. This is the reason the survey was limited to just one day in each area. Despite the drawbacks, the survey provided information on the structure and composition of the groups, the percentage of calves and the proportion of tuskers – information useful to wildlife managers. One clue to the reliability of this method is to repeat the survey immediately. An immediate recount means, as Rodgers (1991) argues, there is no chance of conditions changing. Ideally the survey needs to be carried out twice a year in the dry and rainy seasons to determine the changes in the range and abundance of the elephant.

An index of abundance would be sufficient to tackle many of the wildlife problems. Wildlife management authorities need to know the status of the elephant populations, whether they are increasing, decreasing and whether their numbers should be regulated to reduce conflict and to relax the pressure on habitats (Blanc *et al.*, 2003).


McKay (1973) estimated that there could be between 200-500 elephants in the entire northern Sri Lanka in the late 1960s. The present survey indicates that at least 220 elephants are present in the Mannar District alone, but the numbers could be even as high as 350 or higher. Such information, however imperfect it may be, would be useful for the effective conservation and management of elephants. While the effect of the elephant on its habitat may benefit many other species of wildlife sympatric with it in the area, it can also have a detrimental impact on the vegetation, should elephant density increase.

Mannar District represents an important area for elephants. It provides a connection for the elephants in

Table 2. Estimates of shoulder heights of elephants based on footprint diameter based on the criteria established by Eisenberg and Lockhart (1972) and McKay (1973).

#	fore foot diameter (cm)	circumference (in cm)	shoulder height (m)	size class
1	33	102.3	2.08	adult
2	15	47.1	0.96	calf
3	16	50.3	1.01	calf
4	37	116.2	2.36	adult
5	20	62.8	1.27	juvenile
6	37	116.2	2.36	adult
7	34	106.8	2.17	adult
8	32	100.5	2.04	adult
9	22	69.1	1.40	juvenile
10	28	87.9	1.78	subadult
11	29	92.1	1.85	subadult
12	17	53.4	1.08	calf
13	30	94.2	1.91	subadult
14	35	109.9	2.23	adult
15	18	56.5	1.15	juvenile
16	34	106.8	2.17	adult
17	32	100.5	2.04	adult
18	24	75.4	1.53	subadult
19	32	100.5	2.04	adult
20	35	109.9	2.23	adult
21	30	94.2	1.91	subadult

Number of adults 10 (47.6%), subadults 5 (23.8%), juveniles 3 (14.3% and calves 3 (14.3%)



Artist: Manjula Wijesundara

Wilpattu National Park and those in the Giant's Tank and Madhu Road Sanctuaries. In addition to enhancing elephant movement and gene flow, the area has great value as habitat for plant and animal communities. The survey has shown that elephant distribution is not confined to such protected areas alone. The majority of elephants appear to be found in unprotected areas. Therefore this situation could lead to increased conflict with people, especially in areas where refugees have been resettled and agriculture is the principal land use. There is an opportunity in Mannar District to link up the Wilpattu North Sanctuary with the Giant's Tank and Madhu Road Sanctuaries (Fig. 2) through a forest corridor to enable the elephants to move to and fro across the A14 highway. This is possible given the low human population density in the area and the presence of many water holes and forest blocks. Such a forest corridor provides a means of ensuring connectivity for elephant movement across landscapes threatened with fragmentation. The proposed corridor being more than 10km wide, it would be more than adequate to ensure the gene flow between elephant populations to the north and south of the A14 highway. As Sukumar (2003) argues, if the objective of a corridor is merely to act as a passage for elephants, it would be better to maintain the vegetation in a state that encourages rapid movement rather than a prolonged stay that could promote conflict.

Information on elephant number, minimum viable population (MVP) size, and movement pattern would be invaluable to define clearly the management objectives for both protected and unprotected areas of elephant range (Lindsay, 1993). Furthermore, should a national park, nature reserve or forest corridor be planned in the Mannar District to ensure the long-term survival of the elephant, such information would be most useful.

References

Abayakoon, C.D.F. (1950). Cleopatra's Pearls came from Mannar. *Observer Annual* 107-112

Anon (1988). *Sri Lanka/Swiss Remote Sensing Project, Final Report*. Department of Geography, University of Zurich, and Center for Remote Sensnsing, Survey Department of Sri Lanka, Colombo.

Blanc, J.J., Thouless, C.R., Hart, J., Dublin, H.T., Douglas-Hamilton, I., Craig, C.G. & Barnes, R.F.W. (2003). *African Elephant Status Report 2002: An update from the African Elephant Database*. IUCN/SSC African Elephant Specialist Group. IUCN, Gland, Switzerland and Cambridge, UK.

Boake, W.J.S. 1888. *Mannar – A Monograph*. G.J.A.Skeen. Government Press.

Boyle, D. (1929). Height in elephants. *Journal of the Bombay Natural History Society* **33**: 437.

Brohier, R.L. (1960). Antiquarian Postcripts. No. 26: *Sunday Observer* 25 September (1960)

Caughley, G. (1977). *Analysis of Vertebrate Populations*. John Wiley & Sons, Chichester, UK.

Deraniyagala, P.E.P. (1955). *Some extinct elephants, their relatives, and the two living species*. National Museum, Colombo.

Eisenberg, J.F. & Lockhart, M. (1972). An Ecological Reconnaissance of Wilpattu National Park, *Ceylon. Smithsonian Contributions to Zoology* **101**: 1-118.

Eriyagama, G.J. (1961). The semi-arid vegetation in Mannar region. *The Ceylon Forester* **5**: 66-74

Hednavitharana, W., Dissanayake, S., de Silva, M. & Santiapillai, C. (1994). The survey of elephants in Sri Lanka. *Gajah* **12**: 1-30.

IUCN, (1990). *IUCN Directory of South Asian Protected Areas*. IUCN, Gland, Switzerland and Cambridge, UK.

Jachmann, H. (1996). Direct counts of elephants from the ground. In *Studying Elephants* (ed. K. Kangwana), pp. 49-56. African Wildlife Foundation, Nairobi, Kenya.

Kurt, F. (1969). Some Observations on Ceylon Elephants. *Loris* **11**: 238-243.

Lewis, P. (1894). Archaeology of the Wanni. *Journal of the Ceylon Branch of the Royal Asiatic Society* **13**: 151-176

Lindsay, W.K. (1903). Conservation and management of elephants in Namibia. *Pachyderm* **19**: 49-53.

McKay, G.M. (1973). Behavïor and Ecology of the Asiatic Elephant in Southeastern Ceylon. *Smithsonian Contributions to Zoology* **125**: 1-113.

Moss, C. 1988. *Elephant Memories: Thirteen Years in the Life of an Elephant Family*. Fontana, Williams Collins Sons & Co. London

Nicholas, C.W. (1954a). From the Warden's Field Book: The Ceylon Elephant in Antiquity. (i) The Sinhalese Period. *The Ceylon Forester* **1**: 52-58.

Nicholas, C.W. (1954b). From the Warden's Field Book: The Ceylon Elephant in Antiquity. (ii) The Portuguese, Dutch and British Periods. *The Ceylon Forester* **1**: 103-111.

Nicholas, C.W. (1955). The present status of the Ceylon elephant. *The Ceylon Forester* **2**:38-40.

Nicholas, C.W. (1959). Historical Topography of ancient and medieval Ceylon. *Journal of the Ceylon Branch of the Royal Asiatic Society* **6**: 233 pp.

Norris, C.E. (1959). *Preliminary Report on the Ceylon*

Elephant Field Survey. Wild Life Protection Society, Colombo.

Owen-Smith, R.N. (1988). *Megaherbivores: The influence of very large body size on ecology*. Cambridge University Press, Cambridge.

Ramakrishnan, U., Santosh, J.A. & Sukumar, R. (1991). *Censusing Elephants in Forests*. Asian Elephant Conservation Centre, Bangalore.

Riney, T. (1982). *Study and Management of Large Mammals*. John Wiley & Sons. Chichester, U.K.

Rodgers, W.A. (1991). *Techniques for Wildlife Census in India. A Field Manual*. Wildlife Institute of India, Dehra Dun.

Santiapillai, C. & Wijeyamohan, S. (2003). The impact of civil war on wildlife in Sri Lanka. *Current Science* **84**: 1182-1183.

Shoshani, J., Hagos, Y., Yacob, Y., Ghebrehiwet, M. & Kebrom, E. (2004). Elephants (*Loxodonta africana*) of Zoba Gash Barka, Eritrea: Part 2. Numbers and distribution, ecology and behaviour, and fauna and flora in their ecosystem. *Pachyderm* **26**: 52-68.

Shambaugh, J., Ogelthorpe, J. & Ham, R. (with contributions from Sylvia Tognetti). (2001). *The Trampled Grass: Mitigating the impacts of armed conflict on the environment*. Biodiversity Support Program, Washington D.C.

Sukumar, R., Joshi, N.V. & Krishnamurthy, V. (1988). Growth in the Asian elephant. *Proceedings of the Indian Academy of Science (Animal Science)* **97**: 561-571.

Sukumar, R. (1989). *The Asian Elephant: Ecology and Management*. Cambridge University Press, Cambridge.

Sukumar, R. (2003). *The Living Elephants: Evolutionary Ecology, Behavior, and Conservation*. Oxford University Press, Oxford.

Vane, G. (1887). The Pearl Fisheries of Ceylon. *Journal of the Ceylon Branch of the Royal Asiatic Society* **10**: 14-40

Walsh, P.D., White, L.J.T., Mbina, C., Idiata, D., Mihindou, Y., Maisels, F & Thibault, M. (2001). Estimates of forest elephant abundance: projecting the relationship between precision and effort. *Journal of Applied Biology* **38**: 217-228.

Weerakkody, D.P.M. (2000) *Taprobane: Ancient Sri Lanka as known to Greeks and Romans*. Indicopleustoi archaeologies of the Indian ocean. Brepols, France.

Western, D., Moss, C.J. & Georgiadis, N. (1983). Age estimation and population age structure of elephants from footprint dimensions. *Journal of Wildlife Management* **47**: 1192-1197.

Whitfield, J. (2003). The law of the jungle. *Nature* (London) 421: 8-9.

Wijeyamohan, S., Baheerathi, T., Luxmy, S., Prbha, K., Sajithran, T.M., Sivagini, S., Sivagnanam, V., Theban, S., Wijesundara, C. & Santiapillai, C. (2002). Diversity of birds in the Giant's Tank, Wanni region, Sri Lanka. *Tigerpaper*, **29**: 11-14

Williams, B.K., Nichols, J.D. & Conroy, M.J. (2002). *Analysis and Management of Animal Populations*. Academic Press, London.



Mahouts and their elephants working as Conservation Response Unit in Sumatra

Wahdi Azmi[1], Heidi S. Riddle[2], Agung Setyabudi[3], Wiratno[4], Andi Basrul[5], Edy Sunardi[1], Diding M. Ichsan[1], and Selamat Hardi[1]

The importance of Sumatran Elephants

Due to the large home range and its ecological requirements, the elephant is an ideal flagship and indicator species for the conservation of forest and associated biodiversity in Sumatra. Elephants are also an umbrella species in that they use a variety of habitats occupied by other charismatic species such as Sumatran Rhino, Sumatran Orang Utan and Sumatran Tiger.

Regarding the Asian Elephant, high conservation priorities should be given to populations harbouring the greatest proportion of the species' genetic variability, and to genetically unique populations. Mitochondrial DNA analysis of the Sumatran Elephant has suggested that it represents a separate Evolutionary Significant Unit (ESU), thus supporting their separate sub-specific status. Recent work conducted by Fernando et al. using micro satellite analysis has confirmed the unique ESU status of both the Sumatran Elephant and the Borneo Elephant (Fernando *et al.*, 2003).

Lack of a conservation vision in the land use planning process

As found in other Asian elephant range countries, the fragmentation of elephant habitat in Sumatra is a well-recognised problem. Habitat patches are becoming smaller and are more isolated from each other, increasing the vulnerability of elephant populations. Two thirds of current Asian elephant range is in non-conservation areas (Sukumar, 1989). Most of the current conservation/ protected areas in Sumatra are inadequate in their design to effectively protect Sumatran elephants because they do not protect prime elephant habitat. Most wild elephants are present in blocks of forest outside of conservation/ protected areas, invariably creating a high potential for incidents of human-elephant conflict. These issues reflect the failure of the current land-use planning system to take into account elephant habitat requirements and the long-term benefits of forest conservation. Ramono (2000) stated that the conversion of primary forest into agricultural holdings has been one of the main causes of conservation problems in Sumatra and the elephant has been among the large mammals most seriously affected by it. Development programmes have led to the annual elimination of tens of thousands of hectares of elephant habitat.

Economic incentives have dominated land use planning in Sumatra, resulting in a substantial loss of highly bio diverse forest to cultivation. Most of the lowland forests that are well recognised as containing the richest biodiversity and prime elephant habitat have been allocated for other purposes such as logging concessions, transmigration sites and large-scale plantations. In North Sumatra Province, 900,000 hectares of lowland tropical rain forest has been converted into small-scale and large-scale oil-palm plantation in the period of 1990-2005. This development movement has lead to substantial and higher pressure on Gunung Leuser National Park in the form of land encroachment for small-scale oil plantation.

Despite containing some of Sumatra's most extensive forests only 10% of Aceh is protected as Gunung Leuser National Park and habitat loss continues through legal forest conversion activity, illegal deforestation and settlements both outside and within Protection Forest and Protected Areas. Habitat fragmentation is the by-product of this process and now is threatening to break the chain of forest running along and flanking the central mountainous range of Bukit Barisan from the North in Aceh to the South. As a result, elephant habitat has been scattered and some of the small populations are being isolated in a block of forest, increasing the risk level of extinction. The overlapping of elephant habitat with human activities in the same area also significantly increases the number of human-elephant conflict incidents.

Human-elephant conflict

In the recent past, human-elephant conflict in Sumatra has been handled by capturing the raiding elephants, driving them or scaring them off using sound and fire, while a few plantations use modern technology such as electric fences.

Biological considerations of elephant demography indicate much lower intrinsic rates of population growth in rain forest habitats as compared to drier forests. This means that capturing elephants from rain forests, as a means of containing depredation or for other reasons, is more likely to result in a depletion of the populations as these recover only at a very slow rate. Capture should

Corresponding author
Wahdi Azmi
Fauna and Flora International,
Jl. Garuda No. 61 A Sei Sikambing,
Medan, Indonesia
[1]Fauna and Flora International - Sumatran Elephant Conservation Program;
[2]International Elephant Foundation;
[3]BKSDA Bengkulu,
[4]Gunung Leuser National Park;
[5]BKSDA Aceh.

thus be the last resort of management and should be confined to non-viable populations preferably for translocation, or for identified "rogue" elephants which are a serious threat to human lives (Sukumar, 1999).

The long-term conservation of the elephant in Sumatra, however, requires that elephants and people co-exist with minimal conflict. Otherwise demands for the removal of elephants will be politically difficult to ignore, resulting ultimately in the depletion of elephant populations on the island. Thus we have to find appropriate methods to mitigate elephant-human conflict in Sumatra before the situation reaches a point of no-return for the elephant.

Elephant camps as a result of human-elephant conflict

As a result of escalating human-elephant conflicts in the 1980s, the Indonesian Directorate General of Forest Protection and Nature Conservation (PHKA) ordered the capture of wild problem elephants and Elephant Training Centres (ETCs) were established in six provinces throughout Sumatra. These were later renamed Elephant Conservation Centres (ECCs). In 2001, in collaboration with FFI, PHKA registered 482 elephants in captivity in Sumatra (Janis *et al.*, 2001). The low survival rate in the capturing and training process implies that a higher number of elephants have actually been captured. Most of these elephants are located in government managed Elephant Conservation Centres. An unexpected result of the capturing policy has been over-crowding in all ECCs in Sumatra. To add to this, all the centres lack funds for good management enforcement so camps cannot demonstrate their roles in the global effort of in-situ elephant conservation, which was part of their original purpose.

Establishing Conservation Response Units

The Conservation Response Unit (CRU) concept is founded on the belief that diversity is only secure when diverse conservation strategies are employed. In-situ and ex-situ approaches are not mutually exclusive; no single method of conservation is optimal for all situations, and no single method can succeed alone. Different conservation systems can complement each other and provide insurance against the shortcomings of any one method. Ultimately, the success of both in-situ and ex-situ approaches depends on forging strong links between the two.

The CRU model is but one method that provides a strong link between *in-situ* and *ex-situ* elephant conservation. This model is utilising once neglected captive elephants and their mahouts for direct field based conservation interventions to support the conservation of wild elephants and their habitat, and achieve positive outcomes for both elephants and people. By creating this link, and ensuring that these elephants are seen as an important resource and doing positive deeds, it is expected that local communities, decision-makers and other stakeholders will recognise their contribution and hopefully focus greater attention on protecting Sumatran Elephants, in the wild and in captivity.

This philosophical background of the CRU concept has been positively received by both the national and international elephant community, and the concept is being implemented in a wider part of Sumatra, in a collaboration between Fauna and Flora International (FFI), International Elephant Foundation (IEF) and Balai Konservasi Sumber Daya Alam (BKSDA) as part of the Sumatran Elephant Conservation Strategy with a multiple year partnership commitment.

Local governments and communities should be convinced that Sumatran elephants are worthy assets and genetically important for global elephant conservation. Elephants can play a significant role in generating income for local people through ecotourism and by protecting the function of the forest, which is elephant habitat. Without making this connection and ensuring that these elephants are seen as an important resource, there will be a continuing welfare problem and the important genetic resource that these animals represent (as a high proportion of an ESU of the Asian elephant) will be lost.

Structure and goals of the CRU

The CRU teams are composed of 14 captive elephants from two ECCs (Aceh and Seblat) and 14 of their mahouts, 14 government forest rangers, and 3 FFI conservation officers spread over three CRU posts placed in targeted working areas. Working in partnership with local government, local communities and NGOs, the CRU project has 4 main objectives: 1) mitigating human-elephant conflict; 2) reducing wildlife crime activities in the important elephant habitat through forest patrol and monitoring; 3) raising awareness among local people of the importance of conserving elephants and their habitat; 4) establishing community-based ecotourism to ensure long-term CRU financial sustainability. Working towards field-based conservation intervention, both captive elephants and mahouts have been offered a new alternative future. Captive elephants play an important role by providing transportation during forest monitoring patrol activities, as a tool for gaining local community interest during awareness events, and driving away crop raiding wild elephants should conflict incidents arise. Mahouts, as part of the CRU team, not only take care of the elephants but are involved in all CRU activities and have gained training in wildlife observation techniques and basic use of navigation devices and mapping.

CRU as a site specific project

During the early stages of the CRU operation, various problems and issues specific to each working area were identified and will be the focus of activities in the next phase of the CRU project.

One of the CRU sites, in Bengkulu Province, successfully addresses the land encroachment and illegal logging issue inside the ECC area and identified the need to increase the protection status of the ECC area, as well as expand the protected area and establish a corridor to the Kerinci Seblat National Park. The proposed corridor will increase the protected forest area from 6865 ha. to 18.000 ha. The Bengkulu Nature Conservation Agency (BKSDA) has taken the leading role in proposing this increased protection status to the Indonesian Forestry Ministry, with the endorsement of the local government. Establishing a good management plan for the new proposed protected corridor and elephant camp forest has become the focus of the CRU work in this area.

The CRU in North Sumatra, in collaboration with Gunung Leuser National Park, is working to empower the National Park resort posts in Langkat district through the implementation of CRU activities. Illegal logging and land encroachment still remain the main issues to address. Working closely with the local community, the CRU is also involved in developing eco-tourism in the Tangkahan area, as one way to maintain the CRU objectives by the community itself with support from eco-tourism activities.

In the Tangkahan area, adding revolving positions for three local villagers to the CRU team to learn, assist, as well as provide contributions to all CRU activities on a daily basis offers an opportunity for the CRU team to transfer their skills and share knowledge locally.

Training and capacity building

With the establishment of each CRU team, capacity building has been an initial focus for staff and project partners. Training sessions have been conducted with topics covering:

- Survey and forest monitoring techniques, including basic navigation techniques using a hand held GPS
- Human-elephant conflict mitigation
- Wildlife data recording
- Community awareness

Some CRU staff has been sent to other elephant projects, such as the ones managed by the Wildlife Conservation Society (WCS) to learn about the Way Kambas ECC in Lampung Province, and to share skills with the Crop Protection Unit Team operating under the supervision of WCS.

With the support of IEF, one of the CRU team leaders was sent to the elephant camp at the Jaldapara Wildlife Sanctuary (India) to learn different methods for similar projects (i.e. patrols, camp management, etc). IEF is also committed to support the first mahout workshop in Sumatra to be held in 2006, to establish a communication forum and share knowledge amongst representative mahouts from each of the various camps in Sumatra.

Most of the CRU team members have little educational background, yet through a series of capacity building activities have been trained in assessing and selecting priority areas for CRU activities and field patrols, operating hand held GPS units, filling in standardised data-sheets for forest patrolling and conducting HEC assessments. This empowerment has provided a sense of dignity to the mahouts, a yet unexplored potential source of human resources working for field based conservation.

In the future, with identified field work requirements gathered from experience during implementation of the program, further training needs will be assessed to best approach site specific issues.

Elephant back patrols

Each CRU post is divided into teams who conduct patrols for 7-10 consecutive days a month, during which CRU team members record sightings or evidence of illegal activities, human-wildlife conflicts and wildlife presence. Communities in these critical conservation areas are exposed to elephants in a positive context through their physical presence whilst passing through villages on patrols. The team carries hand-held GPS units to properly identify sighting locations, and a digital camera for documentation purposes. They also fill out report sheets and a narrative when they return from patrol and this data is provided to the partner BKSDA and GLNP offices.

Whilst on patrols, the CRU team records findings as follows:

1. Forest crimes which include illegal logging, land encroachment, illegal hunting, fish bombing, etc.
2. Areas where wild elephants and other wildlife are present, including saltlicks and other means of verification.

Assisted by the data processing group of FFI's Sumatran Elephant Conservation Programme (SECP), information captured on the GPS devices is downloaded and digitally mapped to record patrol routes, illegal activities and wildlife presence in order to best focus patrolling efforts and to explore new areas. Project activities have provided important basic and baseline data about the current extent of forest cover and the status of key biodiversity components in each CRU working area, and this data has been a strong justification for the new forest protection area proposed in Bengkulu-Seblat.

Mitigating human–elephant conflict

The CRU concept addresses human-elephant conflict mitigation not only as an effort to avoid further risk of property loss, but of equal importance is parallel consideration of elephant conservation aspects.

The current CRU teams have been trained and subsequently developed their own capacity to assess HEC mitigation options in their specific working areas. The teams have collected detailed information from field based assessments on various aspects of the issues of conflict.

The pattern of human-elephant conflict, as expected, is intermittent. Teams are responding on a regular basis in the working areas, often driving wild elephants back into the forest and also undertaking detailed assessments of any site damage, especially in the Bengkulu-Seblat area. The presence of the CRU has done much to dispel local fears, and the existence of CRU teams is helping keep the HEC issue under control. The continued presence of the CRU posts will ensure that HEC issues do not create animosity in the local community, which has already led to large-scale elephant killings throughout Sumatra.

Awareness programs

Communities in critical conservation areas are exposed to elephants in a positive context through their physical presence as CRU patrols pass through villages, and as they reduce human-elephant conflicts. These visits are used to reaffirm positive attitudes towards elephants and the link between elephant and habitat conservation, promoting a message of tolerance and understanding of the needs of wild elephants, as well as improving attitudes towards the intrinsic value of wildlife.

Conservation awareness programs conducted by the CRU teams include school visit activities, village visits, slide and film programs, games and competitions for visitors and communities living in surrounding locations. The partner NGOs (FFI and IEF), BKSDA and GLNP have developed flyers with general information about elephants, conservation and the CRU project, as well as similar themed children's booklets which are handed out during these community awareness activities.

Elephant related eco-tourism

Sumatra has a large potential for nature-based tourism to generate income for local communities. Many of Sumatra's megavertebrates, such as elephants, rhinos and tigers, are difficult to view because of their low densities, dense vegetation and difficult terrain. Wild Sumatran elephants, for instance, have rarely been seen by tourists or even researchers. However, at some CRU sites there is the potential to set up elephant related tourism projects. The CRU teams in Tangkahan and Seblat are developing a plan for long-term sustainability, as well as a publicity campaign to promote the issue of eco-tourism.

Conclusion

For Indonesia the CRU concept is intended to be one step in reviewing and adopting a new culture of managing elephants, in the wild and in captivity, with the view to ensure acceptance of improved human-elephant relationships within the framework of the country's development programs.

Acknowledgements

We are grateful to the villagers of Tangkahan for their enthusiasm, to all the field staff for their dedication, to the management of BKSDA and Gunung Leuser National Park for their collaboration and support, and to our donors for financial support.

References

Fernando P., Vidya T.N.C., Ng L.S.G., Schikler P., Melnik D.J. (2003). ConservationGenetic Analysis of the Asian Elephant. *Proceedings of the Symposium on Human–Elephant Relationships and Conflicts, Sri Lanka.*

Ramono W. S. (2001). Integrating Sumatran elephant conservation with development programmes. Paper presented at Workshop on the Management of Elephant Conservation Centres in Sumatra, 9-11 April 2001.

Sukumar R. (1989). *The Asian elephant, ecology and management*. Cambridge University Press Cambridge, UK.

Sukumar R. (1999). Report On Elephant-Human Conflicts In Aceh Province. Fauna & Flora International.

Janis R., Taufik A., Azmi W., Yusuf I., & Suprayogi B. (2001). Registrasi Individu Gajah Pada Pusat Latihan Gajah. PHKA, Jakarta.

The shifting chemical signals of Musth

Heidi S. Riddle, David R. Greenwood, and L.E.L. (Bets) Rasmussen

Studies of captive elephants have significantly contributed to our understanding of how olfaction influences elephant behaviour throughout their lifetime, with special importance during the reproductive years. These studies allow longitudinal sample collections coupled with observations of individual elephants over time – something that is difficult in field conditions. The infusion of data from captive studies continues to fill in details on the physiology of musth and some of this information has been correlated by observations of wild elephant herds (Rasmussen & Perrin, 1999; Rasmussen *et al.*, 2002; Rasmussen *et al.*, 2005; Greenwood *et al.*, 2005).

Elephants receive chemoesthetic sensations, or smell, via several systems. In the main olfactory system the trunk is a conduit to the extensive olfactory epithelium covering the numerous turbinates. The billions of olfactory neurons are sensitive to low levels of gaseous compounds. Throughout the mucosal lining of the trunk are the free nerve endings of the trigeminal system, detecting compounds present in higher, often toxic concentrations. Furthermore, elephants have a third chemodetection system: the vomeronasal organ. It is the interplay between the main olfactory system and the enormous vomeronasal system that gives elephants one of the most sensitive and precise "smell" detection systems among mammals (Rasmussen, 1999).

We first began our chemical communication research on signals released in urine by female Asian elephants during the periovulatory period. Separation techniques of a postulated pheromone were based on the high frequency of flehmen responses by male elephants assessing isolated fractions. This assessment mechanism presumably involves the vomeronasal organ system as the trunk tip places liquids on the opening to the vomeronasal organ ducts in the roof of the mouth. These studies were begun in facilities in the USA and, in the controlled captive environment, led to the identification of a urine-derived acetate and demonstration of the bioactivity of its synthetic form. This female-to-male signal met the criteria of a pheromone (Rasmussen *et al.*, 1996).

What about male-emitted signals - what were these signals and did any facilitate reproductively oriented behaviors? Early studies showed that cyclohexanone in temporal gland secretions of musth elephants elicited behavioral responses from females with calves (Perrin & Rasmussen, 1994). To better understand the nature and function of any signals, we needed to study the chemosensory influences of male elephants as related to the phenomenon of musth.

A serendipitous observation of a young captive Asian male elephant heightened our male-focused chemosensory studies. The observation by one of us that a young teenage Asian elephant in a first musth smelled like honey triggered detailed studies of these young males and their secretions (Rasmussen *et al.*, 2002). Subsequent chemical analyses revealed that young Asian male temporal gland secretions (TGS) were composed of a bouquet of sweet odors: acetates, an alcohol (3-hexen-2-ol) smelling like leaves, and pleasant smelling ketones (acetophenone and 2-heptanone). This finding provided the real impetus for in depth chemosensory studies comparing young and older males, as well as musth and non-musth males.

Not only do young Asian male elephants experiencing their first musth emit sweet compounds, but their behavior is unpredictable and erratic. These teenage musth episodes are short in duration and have been termed "moda" (Chandrasekharan *et al.*, 1992). Older males are not only much larger but our studies have shown their musth has a distinctive chemical signature; mature Asian male elephants in musth secrete a very different mixture of chemicals than teenage males (Rasmussen *et al.*, 2002). The older males are more socially and sexually adept and, importantly, capable of sustaining long periods of musth, sometimes several months in duration. During these extended periods not only do they release secretions distinctive of adult musth, but compounds characteristic of the specific phase of these long musth periods.

The concurrent chemical and hormonal maturation of musth is an integral part of the long process of male maturesence within elephant society. When teenage males reach their early twenties, sweet smelling acetates are no longer detectable in their secretions. Our captive studies have shown that pleasant-smelling compounds are transitionally replaced by carboxylic acids; such acids reduce the pH of the temporal gland secretions as low as pH 5.5. During this transitional phase, trace amounts of an acrid ketal, frontalin [1,5-dimethyl-6,8-dioxabicyclo[3.2.1]octane] a demonstrated pheromone, are occasionally

Heidi S. Riddle
Riddle's Elephant and Wildlife Sanctuary, Post Office Box 715, Greenbrier, Arkansas 72058, USA

Professor David R. Greenwood
School of Biological Sciences, University of Auckland, New Zealand

Professor L. E. L. (Bets) Rasmussen
Department of Environmental & Biomolecular Systems, OGI School of Science & Engineering, OHSU, 20,000 NW Walker Road, Beaverton, Oregon 97006, USA




Fig.1 Proportion of enantiomer forms of frontalin during musth




Fig.2 Frequencies of checks to expelled female urine

O=older males (+30 years); 20s=males + 20 years old; T=teenage males

detected (Rasmussen & Greenwood, 2003; Rasmussen & Greenwood, in prep.). Significantly, we have recently discovered that these young male Asian elephants in musth release both sterically possible enantiomers, or mirror image forms, of frontalin but the proportion of forms varies widely from day to day, with the (+) form usually the more dominant one (Greenwood *et al.*, 2005). However, as males continue to mature, not only do the pleasant odors make way for malodorous ones, but the proportion of mirror image forms stabilizes, especially at mid-musth, to almost 50% (Greenwood et al., 2005). Eventually and particularly during the mid-point of a musth episode, older males broadcast mixtures of less volatile, more alkaline-based, longer-lasting ketones, increasing amounts of the bicyclic ketal, frontalin as an almost racemic mixture of the enantiomers of frontalin (Fig. 1) (Rasmussen *et al.*, 2002; Greenwood *et al.*, 2005).

Based on this observed gradual change of chemical emissions, especially of frontalin, we conducted behavioral bioassays both with whole collected TGS and the synthetic form of frontalin at three facilities with captive Asian elephants in the USA (Riddle's Elephant and Wildlife Sanctuary, the Ringling Center for Elephant Conservation, the Oregon Zoo) and at the Auckland Zoo in New Zealand. Behaviors observed in response to presentations of synthetic racemic frontalin were consistent with the results observed with whole temporal gland secretion from older males whose mid-musth secretions contained almost racemic mixtures of frontalin. These behavioral responses were related to the sex, developmental stage, and physiological status of the responding individual.

Thus frontalin, as a racemic synthetic compound, elicited behavioral responses from both males and females. Female reactivity varied with hormonal state and male reactivity varied with age and musth status. Frontalin, either released in the natural TGS by older males or presented as a synthetic component, attracted reproductively ready females, but elicited apprehension from pregnant females, whereas luteal phase females were indifferent. Among males, the older adult males were mostly indifferent to frontalin in either presentation medium, whereas sub-adult males were highly reactive, often exhibiting repulsion or avoidance. Such differential responses may facilitate the smooth functioning of elephant society. This facilitation may occur among males by clarifying who is in musth, in what phase of a musth episode, and who is not, and also revealing male maturity. Females' differential responses, and thus the impact of chemical signals on them, may also affect their reproductive strategies. Many of these social influences and controls within and between male and female groups, especially those affecting breeding strategies, are mediated in large part through the dual olfactory systems.

Wild Asian male elephants are found in transitory groups if not in pairs or solitary. A male may dramatically increase its home range during the weeks when he comes into musth (Desai & Johnsingh, 1995; Fernando & Lande, 2000). Older males, especially but not exclusively, during the annual musth period apparently maintain a subtle, somewhat loose control over non-musth and younger males, and such control increases in area concurrent with home range expansion during musth. The presence of older, larger musth males is effectively signaled by their distinctive TGS and urine chemical signatures that include specific ketonic components; statistically significant captive studies have demonstrated retreat by young males to both musth males actively secreting or presented whole TGS or selected secreted compounds, such as higher molecular weight ketones and frontalin (Rasmussen & Greenwood, 2003; Perrin *et al.*, 1996). A recent anecdotal observation noted that wild elephants in Assam (NE India) were deterred from coming close to previously regularly raided paddy fields by the presence of an older (+30 years old) captive male tethered near the fields during his musth (Dr. K. K. Sarma, personal communication). Observations of wild herds have noted instances of retreat by young males in moda musth to TGS left on substrates by older males in musth (Dr. V. Krishnamurthy, personal observation). In the wild, both the presence of such older males (in musth or not) and internal physiological conditions of the moda musth state may be decisive in determining a young male's investigative behaviours toward females. Teenage males in moda musth may or may not retreat in the presence of a dominant male (especially one in musth), whereas non-musth teenage males are certain to retreat (Rasmussen *et al.*, 2002).

Captive Asian elephant studies have implicated that from a female perspective, musth males are preferred over non-musth males. Females near ovulation were observed to show more interest in urine from musth males whose serum testosterone levels are above 10 ng/ml and especially those above 20 ng/ml (Schulte & Rasmussen, 1999a). Recently our studies demonstrated the reciprocal - namely that musth males have the most direct access to females and that chemical signals influencing the outcome of male dominance interactions reinforce this superior access by musth males.

We examined this in greater behavioral depth among a wild population of elephants in Asia (Rasmussen *et al.*, 2005). Wild male elephants were identified as individuals and categorized by age (and thus size) and musth characteristics. The quantitative appraisal of musth in each male was provided using attributes of musth that included physiological and behavioural characteristics. These characteristics were scored and combined with the chemical assessment of the presence, absence, or relative amount of selected urinary ketones. These appraisals gave a measure of the maturity and degree of musth in individual wild males when compared to data available from precisely studied captive males. Our field data have shown that musth males, in comparison to non-musth males, interacted much more frequently with pre-ovulatory females and demonstrated a strikingly higher frequency of chemosensory responses to females and their urine (Rasmussen *et al.*, 2005).

Not only do musth males interact with more females and more frequently, but also specific olfactory-related responses and patterns of response vary with age and are different between musth and non-musth males. Adult males (+20 years old) while in musth demonstrate a higher frequency of responses to females than their non-musth counterparts. These responses include both distant and close sniffs, trunk tip checks to both the urogenital region of females and their expelled urine (Fig. 2), and flehmen responses. Musth males also characteristically monitored multiple females in succession. Apparently they are more skilled at detecting by olfaction the precise ovulatory status of females.

Male age (and thus size) also influenced responsivity and eventually access to females. Larger, older males (+30 years old) in musth performed significantly more distant sniffs and total contact chemosensory responses than their younger counterparts in musth and than older males not in musth. Among all three primary contact responses - urogenital checks, expelled urine checks, and flehmens - older males in musth exhibited higher responsivity than younger males in musth, with non-musth young males not showing any contact interactions with females. This avoidance of contact responses toward females suggests that perhaps either experience has taught these young males to avoid direct contact with females during their non-musth period (as females may not be interested), or male behaviours are more rational during non-musth.

Specific premating behaviours (e.g. mounting) were more frequent among young males in musth than the older group, suggesting that experienced older males could judge proximity of females to ovulation more accurately and thus expended less effort on non-fertile females. Further down the age structure, teenage males show much lower frequency of distant sniffs toward females than older males, again suggesting their inexperience in recognizing the distant odor of reproductively active females. These quantitative differences in varied responses among different categories of males offer clues about the biological roles in breeding tactics and elephant social structure.

Age revealed different frequencies of chemosensory responses and premating behaviours when teenage males (both in moda musth and non-musth) were compared with their older counterparts. Older males (+30 years old) exhibited the highest chemosensory responses, but summed chemosensory responses by musth and non-musth teenage males were slightly higher than in young adult (+20 years old) counterparts. This high responsiveness by the maturing teenage male group may reflect attempts by this male category to gain sexual knowledge about females through increased contact chemosensory responses. Inexperienced teenage males may lack knowledge of the chemical identity of the pheromone or the ability to precisely assess concentrations, either of which could be the result of learning. Teenage males are still in an experience-gaining period and thus are deciphering the attributes of female cycles, fine-tuning how to socialize with females, and decoding which males are dominant or in musth or both.

Breath also contains relevant social signals. In the somewhat redundant chemical signaling system of elephants, many communicative compounds are present in the blood and excreted or secreted into the urine, TGS or breath. The focus of much of our efforts has been on the influence of chemical signals and pheromones from the urine and temporal glands on elephant societal interactions (Rasmussen & Krishnamurthy, 2000). Our research has demonstrated a correlation between blood constituents and TGS components, yet many of these same chemical communicator molecules are excreted to the outside environment through the breath. Frontalin, indicative of musth in older males, is an especially redundant signal present in blood, TGS, urine and breath. Over a 10-year period (1994-2004), using special stainless steel evacuated canisters to collect exhalant breath, we analyzed more than 100 samples from 10 captive male elephants – both in musth and non-musth. As Asian male elephants often reduce their caloric intake during the musth state, they lose body condition (Desai & Johnsingh, 1995) and a weight loss of hundreds of kilograms has been documented in captive elephants during a musth period; such weight loss implies first a depletion of fat reserves and eventually muscle breakdown. Our breath sample analyses and physiological measurements in blood confirmed such patterns. Breath volatiles were different between male elephants in musth and those not in musth, i.e. clear-cut qualitative and quantitative differences distinguished that a spectrum of ketones, as well as several related alcohols are significantly elevated (Rasmussen & Riddle, 2004). However, while at first ketone levels increased subsequently pentane, an indication of muscle breakdown, was elevated. This correlates with documented elevations in serum lipase and triglycerides. However, the lack of changes in serum creatinine phosphokinase during musth suggests that muscle metabolism may be less affected than fat metabolism, and only after a lengthy period of starvation (Schulte & Rasmussen, 1999b; Rasmussen & Perrin, 1999).

What makes breath signals special is that they are more ephemeral than urinary or TGS secretions. They are carried in the air, probably in part on aerosols, and are usually rather volatile. They diffuse rapidly in the air medium and are subject to photo oxidation. Thus their message is immediate, often close range and non-persistent, and may be individual-to-individual communication. Further studies on the meaning of chemical signals released in breath may help explain why captive male Asian elephants in musth will blow at each other across fences or under doors, strongly suggesting a transfer of chemical signals between individuals.

In exploring breeding strategies it is noteworthy that the Asian elephant shares some breeding tactics common to other sexually dimorphic cognitive mammals, such as a

roving strategy with similarities to some primates (*e.g.* orangutans) and whales (Singleton & van Schiak, 2002; Whitehead, 1990), while the musth parameter adds a unique feature. Among elephants, highly chemosensory oriented searches aid in successful elephant reproduction. Male elephants are natural roamers and that tendency increases with age and is definitely expanded during musth. Reciprocally, musth influences temporary fusion–fission events as roving males join female groups while tracking preovulatory pheromone concentrations. The pH dependency and the modulation by proteins of the Asian female preovulatory pheromone (Z)-7-dodecenyl acetate) affect its signal lifetime (Rasmussen *et al.*, 2003). Thus, from an olfactory perspective, the two male strategies - roaming and older-controlling-younger - demonstrate a well-balanced, time-dependent system of pheromones. Breeding strategies interwoven into the differential social structure of male and female Asian elephants are apparently influenced by the behavioural and chemical conditions of musth as evidenced by greater chemosensory responsivity and increased mating behaviours by musth males; thus musth plays an important determining factor in reproductive strategies.

How does musth influence reproductive success? This question has been raised in a study of African elephants demonstrating that age, size and musth state interacted to affect a male's reproductive success. Older musth males sired a markedly large number of calves, while non-musth males sired about 10% of the calves in this studied population. Furthermore the genetic data from this study suggests that females mate with more males than previously thought, so male mating behaviour, such as mate guarding, is not a good predictor of paternity as a single musth male does not maintain exclusive access to a female throughout her estrous period (Hollister-Smith, 2005). Paternity studies of Asian elephant populations would be revealing, especially in those with depleted older male populations, to determine reproductive success of younger or non-musth males.

Understanding male elephant societal roles, the functioning of male chemical signals and the subtle shifts in these signals – both in musth and non-musth states - are important for conservation applications.

References

Chandrasekharan, K., Radhakrishnan, K., Cheeran, J., Nair, K. M. & Prabhakaran, T. (1992). In *The Asian Elephant: Ecology, Biology, Disease, Conservation and Management* (E. Silas, M. Nair and G. Nirmalan, Eds.). Kerala Agricultural University, India, pp. 71-74.

Desai, A. A. & Johnsingh, A. J. T. (1995). Social organization and reproductive strategy of the male Asian elephant (*Elephas maximus*) in *A Week with Elephants* (J.C. Daniel & H.S. Datye, Eds.). Bombay Natural History Society, Bombay and Oxford University Press, New Delhi, p. 532

Fernando, P. & Lande, R. (2000). Molecular genetic and behavioural analysis of social organization in the Asian elephant (*Elephas maximus*). *Behav. Ecol. Sociobiol.* **48**, pp. 84-91.

Greenwood, D. R., Comeskey, D., Hunt, M., & Rasmussen, L. E. L. (2005) Chirality in elephant pheromones. *Nature* **438**:1097-98.

Hollister-Smith, J., (2005) Reproductive behavior in male African elephants (*Loxodonta africana*) and the role of musth: a genetic and experimental analysis. PhD Thesis, Department of Biology, Duke University

Perrin, T. E., Rasmussen, L. E. L., Gunawardena R., & Rasmussen R. A. (1996) A method for collection, long-term storage, and bioassay of labile volatile chemosignals. *J. Chem. Ecol.* **21**:207-221.

Perrin, T. E. & Rasmussen, L. E. L. (1994) Chemosensory responses of female Asian elephants (*Elephas maximus*) to cyclohexanone. *J. Chem. Ecol.* 20:2577-2586.

Rasmussen, L. E. L. & Greenwood, D. R. (in prep) Reflections among Asian elephants: chirality counts.

Rasmussen, L. E. L., Krishnamurthy, V., & Sukumar, R. (2005). Behavioral and chemical confirmation of the preovulatory pheromone, Z7-dodecenyl acetate, in wild Asian elephants; its relationship to musth. *Behaviour* **142**: 351-396.

Rasmussen, L. E. L., & Riddle, H. S. (2004) Elephant breath: Clues about health, disease, metabolism and social signals. *JEMA* **15**:24-33

Rasmussen, L. E. L. & Greenwood, D. R. (2003) Frontalin: a Chemical Message of Musth in Asian Elephants. *Chem. Senses* **28**:433-446.

Rasmussen, L. E. L., Lazar, J., & Greenwood, D. R. (2003). The olfactory adventures of elephantine pheromones. *Trans. J. Biochem.* **31**:137-141.

Rasmussen, L. E. L., Riddle, H. S., & Krishnamurthy, V. (2002). Mellifluous Matures to Malodorous in Musth. *Nature* **415**:975-76.

Rasmussen, L. E. L. & Krishnamurthy, V. (2000). How Chemical Signals Integrate Asian Elephant Society: The Known and the Unknown. *Zoo Biol.* **19**: 405-423.

Rasmussen, L. E. L. (1999) Elephant Olfaction. *ChemoSenses* **2**:4-5.

Rasmussen, L. E. L. & Perrin, T. E. (1999) Physiological correlates of musth: lipid metabolites and chemosignal composition. *Physiology & Behavior* **67**: 539-549.


Rasmussen L. E. L., Lee T. D., Roelofs W. L., Zhang A. & Daves, G. Doyle (1996) Insect Pheromone in Elephants. *Nature* **379**:684.

Schulte, B.A. & Rasmussen, L.E. L. (1999a) Signal-receiver interplay in the communication of male condition by Asian elephants. *Anim. Behav.* **57**: 1265-1274.

Schulte, B. A. & Rasmussen, L.E.L. (1999b) Ecological and biochemical constraints on pheromonal signaling systems in Asian elephants and their evolutionary implications. *Advances in Chemical Communication in Vertebrates 8* (eds. R. E. Johnston, D. Muller-Schwarze & P. W. Sorenson) Kluwer/Academic/ Plenum Press, pp. 49-62.

Singleton, I. & Van Schaik, C. P. (2002) The social organization of a population of Sumatran Orang-utans. *Folia Primatol.* **22**:877-911.

Whitehead, H. (1990) Rules for roving males. *J. Theor. Biol.* **145**:355-368.



Artist: Manjula Wijesundara

Lipid profile of captive Sri Lankan elephants

W.D. Ratnasooriya, G.K.H.de Alwis, R.D. Wijesekara, R.M. Amarasinghe, and D. Perera

Abstract The lipid profile of the Sri Lankan elephant (*Elephas maximus maximus*) is hitherto not scientifically documented. The aim of this study was to determine the normal lipid profile of adult captive Sri Lankan elephants. Blood was obtained from 78 individuals and lipid levels were determined using enzymatic techniques. The results show that the mean total serum cholesterol level was 44.28±15.52 mg/dl (mean ±SD) (males: 46.06±13.59 mg/dl, females: 42.83 ± 13.44 mg/dl), HDL-cholesterol was 41.49±9.19 mg/dl (males: 44.94±8.05 mg/dl, females: 38.54±9.33 mg/dl) and triglycerides was 25.28 ± 10.69 mg/dl (males: 25.49 ± 8.54 mg/dl, females: 25.14 ± 11.99 mg/dl) the mean total cholesterol / HDL cholesterol ratio was 1.22±0.36 (males: 1.17±0.42, females: 1.28±0.28). LDL-cholesterol was not detectable and therefore LDL/HDL-cholesterol ratio could not be computed. Further, there was no significant gender difference amongst these parameters. This is the first study to record base line data on lipid profile of Sri Lankan elephants.

Key words: *Elephas maximus maximus* Sri Lankan elephant, lipid profile, cholesterol, HDL-cholesterol, LDL-cholesterol. triglycerides

Introduction

Very few studies have been conducted on the haematology, blood biochemistry or genotyping (based on blood) of *Elephas maximus maximus*, the elephant of Sri Lanka, which is critically endangered and included in the Appendix I of the Convention of International Trade on Endangered Species of wild Fauna and Flora (CITES). Regrettably, currently, there are only between 3000-4000 wild elephants and 186 captive elephants in Sri Lanka (Kurt & Mar, 2003). These studies include investigations of blood counts (Ratnasooriya *et al.*, 1990; Ratnasooriya *et al.*, 1993; Silva & Kuruwita, 1993a; Silva & Kuruwita 1993b), osmotic fragility of red blood cells (Silva & Kuruwita, 1994), blood hormonal levels (Ratnasooriya *et al.*, 1992; Ratnasooriya *et al.*, 1993; Lincolin & Ratnasooriya 1996; Poole *et al.*, 1997), blood levels of some ions (Kuruwita, 1993a; Silva & Kuruwita 1993b), enzymes (Kuruwita, 1993a; Silva & Kuruwita 1993b), glucose (Ratnasooriya *et al.*, 1999), triglycerides (Kuruwita, 1993a; Silva & Kuruwita 1993b), proteins (Kuruwita, 1993a; Silva & Kuruwita 1993b), total cholesterol (Ratnasooriya *et al.*, 1995) and mitochondrial haplotypes (Vandebona *et al.*, 2002). However, as yet, a full blood lipid profile (including HDL- cholesterol and LDL-cholesterol) of the Sri Lanka elephant is not scientifically documented. Such data is quite useful directly in diagnosis, treatment, breeding and general welfare and indirectly in the long term conservation and sound management of our elephant. In addition, zoologically, documentation of whatever data on the Sri Lankan elephant is important in view of its endangered status. The aim of this study was to investigate the serum lipid profile of Sri Lankan elephants. This was done on elephants brought to Colombo from various parts of the country to participate in the Navam Perahara (a cultural pageant) in February 1997 and 1998 and elephants from the Pinnawala elephant orphanage.

Prof. W. D. Ratnasooriya
Department of Zoology
University of Colombo,
Sri Lanka.

G.K.H.de Alwis, R.D. Wijesekara, R.M. Amarasinghe, D. Perera
Department of Chemistry
University of Colombo,
Colombo 3,
Sri Lanka.

Methods

A total of 78 adult (35 males and 43 females), apparently healthy elephants, who participated in the Navam Perahera, in 1997 and 1998, and Pinnawela elephant orphanage were the subjects of this study. No attempt was made to differentiate these elephants into the five morphotypes.

Blood samples (5-10 ml) were collected (in standing position) from a vein or artery on the posterior side of either ear (using aseptic precautions) without using a sedation, (between 7.00 h - 12.00 h), using a butterfly needle (18 gause) connected to a plastic 10 ml syringe. The entire bleeding procedure lasted 1.0 - 1.5 min.

Blood was allowed to clot at room temperature (28 - 31°C) and the serum was separated within three hours of collection by centrifugation at 500g for 20 min. The serum was stored at -70°C until the total serum lipid profiles were made. Total serum cholesterol was assayed by an enzymatic procedure using a commercial reagent kit (Randox, Ireland). In this procedure cholesterol esters are hydrolysed by cholesterol esterase to cholesterol and fatty acids. The cholesterol is oxidized by cholesterol oxidase to cholestene3-one and hydrogen peroxide. The

Table 1. Mean (±SD) of serum lipid and lipoprotein levels of adult Sri Lankan elephant (Elephas maximus maximus). Ranges are giving in parenthesis.

Parameter	Entire Group	Males	Females
Total Cholesterol (mg/dl)	44.28 ± 15.52 (28.20 – 85.88)	46.06 ± 13.59 (18.20 – 85.88 ≠)	42.83 ± 13.44 (18.20 – 73.14)
HDL – Cholesterol (mg/dl)	41.49 ± 9.19 (23.45 – 58.65)	44.94 ± 8.05 (28.59 – 58.65)	38.54 ± 9.33 (23.45 – 53.60)
LDL – Cholesterol (mg/dl)	Non Detectable	Non Detectable	Non Detectable
Total Cholesterol / HDL – Cholesterol (mg/dl)	1.22 ± 0.36 (0.68 – 2.01)	1.17 ± 0.42 (0.68 – 2.01)	1.28 ± 0.28 (0.98 – 1.69)
LDL / HDL Cholesterol (mg/dl)	Cannot be Computed	Cannot be Computed	Cannot be Computed
Triglycerides (mg/dl)	25.28 ± 10.69 (12.24 – 55.90)	25.49 ± 8.54 (12.24 – 47.16)	25.14 ± 11.99 (11.19 – 55.90)

Table 2. The mode, median and variance of total cholesterol, HDL cholesterol and LDL-cholesterol in the serum of the Sri Lankan (Elephas maximus maximus) population sampled.

		Mode	Median	Variance
Total cholesterol	Male	-	42.63	184.75
	Female	-	40.08	180.76
	Entire group	62.45	41.92	182.78
HDL cholesterol	Male	-	45.75	64.82
	Female	37.44	38.15	87.07
	Entire group	37.44	42.63	84.38
Triglyceride	Male	-	26.63	72.86
	Female	11.64	21.68	143.86
	Entire group	47.16	24.33	114.37

chromophore quinoneimine is formed from hydrogen peroxide and 4-aminoantipyrine in the presence of phenol and peroxidase (Richmond, 1973). Ten microliters of the serum sample or standard cholesterol solution (200 mg/dl) were incubated with 1000 microliters of the reagent mix at room temperature (28°C) for a period not less than 10 minutes. Absorbance was measured at 500 nm using a Shimadsu double beam spectrophotometer (UV – 21005, Schimadzu Crop., Kyoto, Japan) against a reagent blank. The readings were completed within one hour of incubation. The test is linear up to a cholesterol concentration of 750 mg/dl.

HDL-cholesterol was determined using Randox Test kits. In this procedure low density lipoproteins [LDL (low density lipoprotein) & VLDL (very low density lipoprotein)] and chylomicron fraction are precipitated quantitatively by the addition of phosphotungstic acid in the presence of magnesium ion. After centrifugation, the cholesterol concentration in the HDL (high density lipoprotein) fraction, which remains in the supernatant was determined as follows. The serum sample (200 μl) was mixed with the precipitant (500 μl) and allowed to sit for 10 minutes at room temperature. The mixture was centrifuged for 10 minutes at 4,000 rpm. and the clear supernatant was separated within two hours. The HDL-cholesterol content in the supernatant was determined by mixing the supernatant (100μl) with the reagent (1000μl), incubating for 5 minutes at 37°C, and measuring the absorbance of the sample (A_{sample}) and standard ($A_{standard}$) at 500 nm against the reagent blank within 60 minutes. (The Concentration of HDL cholesterol in the supernatant = A_{sample}/ A_{std} x concentration of standard.) LDL-cholesterol was determined using the formula:

LDL cholesterol (mg/dl) = Total cholesterol (mg/dl) - Triglycerides 5 (mg/dl) - (HDL - cholesterol) (mg/dl).

The serum triglycerides are determined after enzymatic hydrolysis with lipases. The indicator is a quinoneimine formed from hydrogen peroxide, 4-aminophenazone and 4-chlorophenol under the catalytic influence of peroxidase. The sample was mixed (10μl) with the reagent (1000μl) and incubated for 5 minutes at 37°C. The absorbance of the sample (A_{sample}) and standard ($A_{standard}$) were measured at 500 nm, against the reagent blank within 60 minutes.

[Triglyceride concentration = (A_{sample}/ $A_{standard}$) x 200 (mg/dl)]

The test is linear up to a triglyceride concentration of 150 mg / dl. Two preassyed quality control sera (Randox, Ireland) were used as positive controls to monitor accurately. The results are represented as means ± SD. Statistical analysis were made using Mann-Whitney U-test. Significance was set at $P < 0.05$.

Results

All the serum samples made in this study were free of any marked haemolysis and of yellow colouration on visual examination. The results of the blood lipid parameters monitored or computed are summarized in Tables 1 and 2. LDL-cholesterol was either absent or found below the detectable level of the assay kit used. Although, the males generally had a slightly higher level of the individual parameters monitored, none was significantly different ($P > 0.05$).

Discussion

This study records for the first time the random lipid level profile of captive Sri Lankan elephant (*Elephas maximus maximus*). Sequential and fasting blood samples were not collected due to logistic problems. However, according to some, the lipid profile following a meal may be a more important indicator of coronary heart disease risk than the fasting level (Ryu *et al.*, 1992). Further, we have previously determined blood glucose (Ratnasooriya *et al.*, 1999) and cholesterol levels (Ratnasooriya *et al.*, 1995) of Sri Lankan elephants under random conditions rather than in fasting states. Blood samples were collected from apparently healthy animals always in a single day, in a standing position between 7.00-12.00 hours, without applying any pressure to the blood vessels of the ear to minimize variations in lipid profile due to posture, time of the day, and physical state or stress (Dart *et al.*, 1990). Lipid profiles were monitored using enzymatic procedures which are claimed to be sensitive and reliable (Richmond, 1973) and which are widely used in humans. In this study, the lipid profile was determined in 78 individuals, which is a sizable number to provide meaningful data: the current number of captive elephants in Sri Lanka is reported to be 186 (Kurt & Mar, 2003). Collectively, these procedures used allows the data obtained to be regarded as representative and to be considered as reference base line data for captive Sri Lankan elephants as all these captive elephant had their origin in the jungles. It is generally recognized that for each country a base line data on any biological parameter is a must. The overall total cholesterol level reported in this study is in close agreement to what has been reported by us previously using a lesser number of animals (Ratnasooriya *et al.*, 1995).

Both the blood cholesterol and triglyceride level were low in comparison with humans an omnivore (Hahn & Payne, 1997) and presumably in carnivores. This hypolipdaemia is likely to be related to the herbivorous diet of the elephant. Captive Sri Lankan elephants are mostly fed with a monotonous menu: logs of *Caryota urenus* Linn., fronds of *Cocus nucifera* Linn. and leaves and twigs of *Artocarpus heterophyllus* Linn. (Illangakoon., 1993; Godagama, 1999) which are rich in both soluble (pectins, gum and mucilage) and insoluble (cellulose, lignin and insoluble non cellulose polysaccarides) fibre. Soluble fibre is known to induce hypolipidaemia (Bhattacharya & Bhattacharya, 1994) and this may account for low levels of cholesterol and triglycerides in elephants. Increase clearance of these may also precipitate hypolipidaemia.

The most striking observation in this study is the undetectable level of blood LDL-cholesterol (bad cholesterol) and desirable level of HDL-cholesterol (good cholesterol). Lack of LDL-cholesterol is yet another possible cause for a low level of blood cholesterol as it is the main carrier of cholesterol in the blood serum (Holum., 1990; Vance & Van den Bosch., 2000). A high level of HDL and low level of LDL-cholesterol makes the elephants less prone to atherosclerosis and eventually lowers risk of cardiovascular diseases and stroke. This notion is supported by the low cholesterol: HDL ratio and zero LDL: HDL ratio, which are considered as atherogenic indices (Loke *et al.*, 1991; Agarawal *et al.*, 1998), evident in this study. Interestingly, high levels of LDL and low levels of HDL-cholesterol is positively linked with arterial plaque formation (Holum., 1990; Vance & Van den Bosch, 2000) and are reported to be protective against astherosclerosis and cardiovascular diseases (Holum, 1990; Vance & Van den Bosch., 2000). High HDL level in elephants may be due to a high fiber diet, exercise (as several were working elephants) or due to presence of a metabolite/s which acts as fibrate group of lipid regulating drugs (Anon, 2000). Alternatively, it could be due to some hereditary influence, as in humans, whose high HDL-cholesterol levels are determined partially by inheritance (Holum, 1990).

Similarly, several mechanisms may be responsible for the undetectable LDL-cholesterol level in elephants. Presence of a large number of LDL receptors in the liver and/or having high affinity LDL receptors to cholesterol (Vance & Van den Bosh., 2000) in the liver are two striking possibilities: Patient with familial hypercholesterolemia have a defect in the function of LDL receptor and/or low LDL receptors (Holum, 1990). Enhanced physical activity is another mechanism (Hahn & Payn., 1997). Alternatively, metabolites may be present in the elephant liver which may competitively inhibit 3-hydroxy-3-methylglutrayl coenzyme-A (HMG Co-A) as statins, a class of potant lipid lowering drugs (Anon, 2000). Obviously, additional experiments are required to elucidate these potential mechanisms. Understaking such studies is important as it could lead to development of new lipid regulating drugs.

In this study, of the lipid parameters monitored or computed, a significant difference was not evident between males and females. In contrast, in a previous study of ours (Ratnasooriya *et al.*, 1995) a slight but significantly higher level of cholesterol was evident in male elephants. This could be due to the small sample size of the previous study. In conclusion, this study reports for the first time the lipid profile of the Sri Lankan elephant.

Acknowledgments

This study was supported by a grant from National Science Foundation of Sri Lanka (RG/95/C/06). Thanks are also due to Dr. D.S. Kodikara and other veterinary sergeants at Pinnawala elephant orphanage for collecting the blood samples.

Reference

Agarwal, C.G., Verma, R., Manglik, A.K., Ghatak, A. & Kapoor N.K. (1998) High density lipoprotein-Cholesterol in non insulin dependent diabetes mellitus. *Indian Medical Gazette,* **2:** 47-50.

Anon. (2000) *British National Formulary* British Medical Association and Royal Pharmaceutical Society of Great Britain, BMJ Books, London.

Bhattacharya, K. & Bhattacharya, S.K. (1994) Effects of psyllium husk (NATUROLAX) in hyperlipidaemia in clinical study. *Antiseptic*, **I:** 252-256.

Dart, A.M., Reid, C., Fennings, G.L., Conyers, R.A.F. & Nicholls, E.M. (1990) Interpretation and significance of a high blood cholesterol. *Australian Prescriber*, **13:** 81-83.

Godagama, W.K., Wemmer, C. & Ratnasooriya, W.D. (1999) The diet of Sri Lankan domesticated elephant (*Elephas maximus maxmus*). *Vidyodaya Journal of Science*, **8:** 75-84.

Hahn, D.B. & Payne, W.A. (1997) *Focus on health*, WCB McGraw-Hill, New York.

Holum, J.R. (1990) *Fundamentals of General, Organic and Biological chemistry,* Jhon Wiley and Sons, New York.

Illangakoon, A. (1993) A preliminary study of the captive elephants in Sri Lanka. *Gajah*, **11:** 29-42.

Kurt, F. & Mar, K.U. (2003) Guidlines for the management of captive Asian elephants and the posible role of the IUCN/SSC Asian elephant specialist group. *Gajah*, **22:** 30-42.

Lincolin, G.A. & Ratnasooriya, W.D. (1996) Testosterone secretion, mustl behaviour and social dominance in captive male Asian elephants living near the equator. *Journal of Reproduction and fertility.* **108:** 107-113.

Loke, D.F.M., Viegas, O.A.C., Kek, L.P., Rauff, M., Thai, A.C. & Ratnam, S.S. (1991) Lipid profiles during and after normal pregnancy. *Gynecology and Obstetrics Investigation,* **32:** 144-147.

Poole, T.B., Taylor, V.J., Fernando, S.B.U., Ratnasooriya, W.D., Ratnayake, A., Linconl, G., Maneilly, A., & Manatunga, A.M.V.R. (1997) Social behaviour and breeding physiology of a group Asian elephants. *International Zoo year Book.* **35:** 297-310.

Ratnasooriya, W.D., Amarasinghe, A.B.C. & Kodikara, D.S. (1995) Total serum cholesterol levels of Sri Lankan elephants (*Elephas maximus maxmus*). *Ceylon Journal of Science*, **24:** 11-16.

Ratnasooriya, W.D., Caldera, H.S., Premakumara, G.A.S., Liyanage, G.K., Manatunga, A.M.V.R. &

Fernando, S.B.U. (1993). Some haematological values during normal pregnancy in the Sri Lankan elephant (*Elephas maximus maxmus*). *Medical Science Research*, **21:** 153-156.

Ratnasooriya, W.D., Fernando, S.B.U. & Manatunga, A.M.V.R. (1992) Serum testosteron levels of Sri Lankan female elephants (*Elephas maximus maxmus*). *Medical Science Research*, **20:** 79-80.

Ratnasooriya, W.D., Fernando, S.B.U. & Manatunga, A.M.V.R. (1993) Serum prolactin levels of Sri Lankan elephants (*Elephas maximus maxmus*). *Medical Science Research*, **21:** 259-261.

Ratnasooriya, W.D., Fernando, S.B.U. Manatunga, A.M.V.R., Caldera, H.S., Liyanage, G.K. & Premakumara, G.A.S. (1990) Haematological values for adult Asian elephants (*Elephas maximus maxmus*) at the Pinnawela elephant orphanage, Sri Lanka. *Medical Science Research*, **18:** 899-902.

Ratnasooriya, W.D., Gunasekera, M.B., Goonesekera, N.C.N., Vandebona, H.& Kodikara, D.S. (1999) Serum glucose levels in captive Sri Lankan elephants (*Elephas maximus maxmus*). *Vidyodaya Journal of Science*, **8:** 127-134.

Ratnasooriya, W.D., Liyanage, G.K., Caldera, H.S., Premakumara, G.A.S., Manatunga, A.M.V.R. & Fernando, S.B.U. (1992) Thrombocytosis: a possible signal of conception in elephants ? *Medical Science Research*, **20:** 801-802.

Richmond, W. (1973) Preparation and properties of a cholesterol oxidase from *Nocardia sp.* and its application to the enzymatic assay of total cholesterol in serum. *Clinical Chemistry*, **19:** 1350-1356.

Ryu, J.E., Howerd, G. & Craven, T.E. (1992) Postprandial triglyceridemia and carotid atherosclerosis in middle-aged subjects. *Stroke*, **23:** 823-828.


Silva, I.D. & Kuruwita, V.Y. (1993a) Haematology, plasma and serum biochemistry values in free-ranging elephants (*Elephas maximus celanicus*) in Sri Lanka. *Journal of Zoo and Wildlife Medicine*, **24:** 434-439.

Silva, I.D. & Kuruwita, V.Y. (1993b) Haematology, plasma and serum biochemistry values in domesticated elephants (*Elephas maximus ceylanicus*) in Sri Lanka. *Journal of Zoo and Wildlife Medicine*, **24:** 440-444.

Silva, I.D. & Kuruwita, V.Y. (1994) The osmotic fragility of erythrocytes of the Asian elephants (*Elephas maximus*). *Gajah*, **13:** 25-29.

Vance, D.E. & Van den Bosch, H. (2000) Cholesterol in the 2000. *Biochimica Et Biophysica ACTA*, **1529:** 1-8.

Vandebona, H., Goonesekera, N.C.W., Tiedemann, R., Ratnasooriya, W.D. & Gunasekera M.B. (2002) sequence variation at two mitochondrial genes in the Asian elephant (*Elephas maximus*) population of Sri Lanka. *Mammalian Biology*, **67:** 193-205.



Artist: Manjula Wijesundara



Photo: Heidi Riddle for International Elephant Foundation (IEF)

Short Communication

Information on elephant keepers in Sri Lanka

A. Dangolla and I. D. Silva

Abstract Approximately 200 domesticated and privately owned elephants play a significant role in cultural, religious and state functions, and in tourism in Sri Lanka. Information on elephant keepers is vital in order to educate them on better welfare, health care and management of elephants. This aspect has not been scientifically investigated in Sri Lanka in the past. Therefore, data on 53 elephant keepers were collected by a detailed pre-tested questionnaire administered during personal interviews with them during August 2001-June 2002. All keepers were Sinhala (race), Buddhist (religion), males; 49% were over 41 years of age. A majority had studied up to grade 8 (79%), and could write Sinhala language comprehensively (72%) and could read Sinhala news papers (83%). They provided satisfactory level of education to their children. A majority (70%) were receiving a monthly salary of more than that of an average laborer in Sri Lanka. Elephants in the tourist trade worked almost 30 days/month. Most logging elephants worked only 4-5 hours a day. Food was not provided for the elephants while working, by 58% of the keepers and most keepers provided water to their elephants during work (89%). Most of them did not provide sufficient food immediately after work (79%) though food was provided after allowing the elephant to bathe, immediately after work. Twenty two (41%) keepers had changed their assistants repeatedly. Most keepers (75%) carry the goad (5-6 feet long wooden pole with a steel hook at the end used for restraint), a stick and the knife all the time. Almost all keepers (91%) knew only about herbal medicaments for different ailments in elephants, probably because they had either no faith on, or had little or no exposure to western medicines both for their elephants as well as on themselves. Approximately half (47%) consumed alcohol daily. Thirty percent (30%) of the keepers smoked between 1-5 cigarettes daily and 98% chew beetle leaves. **Keywords:** Elephant, Elephant keeper

Introduction

At present, there are approximately 200 elephants (both domesticated and privately owned) in Sri Lanka, and the domesticated elephants are distributed in 12 out of 25 districts in the country. This number is similar to that of privately owned Asian elephants in USA (Roocroft & Atwellzol, 1994). The Sri Lankan elephants walk or at times travel in trucks, over long distances during annual festival seasons to participate in cultural, religious and state functions throughout the country. They also play a vital role in generating income from the developing tourist industry. A rapid decline in the future population is anticipated since more than half of the elephants are over 50 years old and therefore not in breedable ages. The net result of this reduction may adversely affect the economy and the people of Sri Lanka in many ways.

The Captive Elephant Owners Association of Sri Lanka (CEOASL) has taken several vital steps to improve the management of these elephants. The elephant owners hire keepers (all males) to manage their elephants. Elephants therefore, spend most of their time with the keepers. The CEOASL has decided to collect information on elephant keepers to find the ways in which they work the elephants, in order to find ways and means of educating and training them on better management and health care practices with the objective of improving the welfare aspects in the hope of sustaining the existing elephant population. Such data on elephant keepers has not been documented in Sri Lanka. Each domesticated elephant has one keeper and one or two assistants to the keeper.

A. Dangolla and I. D. Silva
Department of Veterinary Clinical Sciences,
Faculty of Veterinary Medicine and Animal Science,
University of Peradeniya,
Sri Lanka.
Email: adangolla@yahoo.com

Materials and methods

Information of elephant keepers in 10 districts in which majority of the domesticated elephants live, were collected via a questionnaire which constituted of aspects on keeper's family, education of the keeper and his children, ability to read and write, his income, whether he smokes and consume alcohol, working schedule of the elephant and methods he use to restraint elephants. After pre-testing the questionnaire, necessary changes were made and the final version was administered via personnel interviews to 53 keepers during August 2001 to June 2002.

Results

Family, Income and level of education

The race and religion of all elephant keepers interviewed were Sinhala (race) and buddhists (religion) respectively, by birth. Nearly a half (26; 49%), were over 41 years old and they reported to have a maximum of two children. Only one keeper had studied up to grade 12 (General

Certificate of Education, Advanced level), and 42 (79%) only up to grade 8. Thirty-eight (72%) were literate in the Sinhala language and 44 (83%) could read Sinhala newspapers. All of their school-going-age children were attending school. None of the children had left formal education at an early age. Eighteen (34%) keepers indicated their monthly income to be less than Rs. 4000.00 and 37 (70%) indicated a monthly income of above Rs. 5000.00.

Working the elephant

Twenty-two (40%) keepers worked the elephants under their care, up to an average of 10 days a month. However, those elephants in the tourist trade, especially the aged females, work almost 30 days a month. Their work, consist of walking approximately 5 kilometers per day while carrying 2-4 tourists.

Thirty four (64%) keepers assist the owners to decide on the work place. Most keepers had the habit of visiting the workplace before commencing the work. Those elephants that were hired for logging, which is strenuous, work only for 4-5 hours a day. However, 6 (11%) keepers worked their elephants up to 10 hours a day, at times. Twenty-two (41%) keepers changed their assistants repeatedly for some reason.

Providing food and water to the elephant

Food was not provided for the elephants during work by 31 (58%) keepers, and 47 (89%) provided water to their elephants during work, 42 (79%) provided adequate drinking water immediately after work and the food was given subsequently. In general, they were bathed immediately after work for about 4 hours a day, and most of the food was given thereafter.

General

Majority of keepers (40; 75%) carry the goad (5-6 feet long wooden pole with a steel hook at the end), a stick and a knife at all times, but they mainly use the stick to control the elephant. Almost all keepers (48; 91%) knew a variety of minor herbal medicaments used for different ailments of elephants. Only 4 keepers were aware of western anthelmintic preparations used on elephants, while all of them were confident and had used ayurvedic medicaments for various ailments in elephants under their care.

Approximately half (25; 47%) of the keepers interviewed consumed alcoholic liquor, 30% smoked cigarettes and 98% chewed beetles. Beetle leaves are traditionally chewed with areca nut, tobacco and lime. One third of keepers (17; 32%) chew beetles at least 17 times daily.

Discussion

It was not surprising that all keepers interviewed are Sinhala buddhist males, as most functions in which elephants participate are associated with Buddhist viharas or Hindu temples. The present culture in Sri Lanka does not appear to attract and favor female keepers, probably because of the physical hardships of this occupation. Though most keepers had not completed their primary education, they could read and write reasonably well and they appeared to be very concerned over the education of their children. Most keepers had left school education early and had learnt the trade of elephant keeping as a profession from their fathers, grand fathers or uncles, as they were fascinated about elephants. It is likely that the number of professional elephant keepers who would learn the profession from their fathers would be reduced and therefore, a formal training of keepers may become inevitable in future. Such training would also provide an opportunity to educate them on welfare aspects, newly reported diseases of Sri Lankan elephants (Dangolla *et al.*, 2002) and also other health risks due to excessive work load (Silva & Dangolla, 2002).

Most elephant keepers dress shabbily though their income is more than that of an average laborer. It is also known that these keepers earn incomes in addition to their regular pay unless the elephant falls sick for long periods of time. The keepers of male elephants face financial difficulties when their elephants come in to "musth" as they are traditionally not worked during this period.

Examination of the place of work by most keepers prior to commencement of work is commendable since a joint decision could then be made with the owner. This is important since several elephants have died due to falls when working in bad terrains. The practice of not offering even small amounts of food while at work may not be advisable considering the relative inefficiency of the digestive tract of elephants.

The goad and the stick carried by the keepers were for restraint while the knife was for cutting the branches and leaves for food and to trim the nails of elephants while bathing, when necessary. However, there had been many instances of long term wound dressing and antibiotic therapy on injuries caused by keepers using sharp goads indiscriminately.

Many elephant keepers had not used western medicines on the elephants under their care possibly because they (both elephant and themselves) have had little or no exposure to western medicine or because of lack of faith. We were able to convince many keepers that regular administration of a haematinics to old elephants, especially those in tourist trade, could sustain good health status (Dangolla & Silva, 2000).

The respondents appeared to be quite addicted to chewing beetle leaves and consuming alcoholic liquor. Interestingly, elephant keepers are known to tell "untruths" (Evans, 1961). However, the fact that most of them had indicated a reasonably higher income may suggest that the extent of lying was possibly less.

It is apparent from the work load, the diet and the diseases emerging in elephants, that the keepers need to be educated promptly on those facts, if the older domesticated elephants are to be preserved longer.

Acknowledgements

D Weerasiri and ULD Jayantha are thanked for their assistance during the study.

References

Dangolla, A. & Silva, I..D (2000). A case of mal-absorption syndrome due to fractured molars in an Asian elephant (*Elephas maximus maximus*). *Sri Lanka Vet. Journal*, 47(1A):11-12.

Dangolla, A., Silva, I., Thevenasan, V., Siribaddana, A., Ratnathunga, N., Perera, J. & Fernandopulle, S. (2002). Preliminary information on Tuberculosis in an Elephant (*Elephas maximus maximus*). *Proceedings of 54th Annual Convention of the Sri Lanka Veterinary Association*, 3-4th May 2002, Gannoruwa, Sri Lanka

Evans, G.H. (1961). *Elephants and their diseases: A treatise on elephants.* Supdt. Govt. printing and staty., Union of Burma, Rangoon. 323 pp.

Roocroft, A. & Atwellzoll, D. (1994). *Managing elephants; an introduction to their training and management.* Fever Tree Press, PO Box 205, Ramona, Ca 92065. USA. 198 pp.

Silva, I. & Dangolla, A. (2002). Blood levels of cholesterol and triglycerides in wild and domesticated Asian elephants (*Elephas maximus maximus*) in Sri Lanka.



Photo: Heidi Riddle for International Elephant Foundation (IEF)




Photo: Heidi Riddle for International Elephant Foundation (IEF)

Short Communication

Elephant Endotheliotropic Herpes Virus (EEHV)


Charlie Gray

Elephant Endotheliotropic Herpes Virus (EEHV) is a disease that has been the cause of death of many otherwise healthy Asian elephants. It was first identified as the cause of death in a 16 month old Asian elephant at the National Zoo in Washington, D.C., USA. To date there have been over 30 fatal cases in North America and Europe, the majority of the fatalities have been young animals. In Europe, losses are reported as 25 % of their newborn Asian elephant populations due to EEHV, and 20 % of Asian elephants under the age of ten years are lost to EEHV in North America. This disease is a serious threat to free-ranging as well as captive elephants. Recent information indicates that EEHV is also found in captive and wild populations of Asian elephants in range countries. EEHV and the disease it causes are very complex and require a great deal more research.

There are two types of disease presentation associated with EEHV, acute with the onset in days to hours and peracute or sudden death. Early symptoms of the disease can include bloodshot eyes, head and neck swelling, lethargy, cyanosis of the tongue and ulcers in the hard palate of the mouth. EEHV infects the micro vascular endothelial cells of the heart, liver and tongue. Viral-induced endothelial injury leads to critical hemorrhages with death attributed to cardiac failure. Transmission of EEHV is unknown and the virus is very difficult to culture. There have been four elephants with EEHV that have survived the disease all were treated immediately after diagnosis with the anti-viral drug famciclovir

In September of 2005 more than 20 animal health specialists from around the world met in Houston, Texas, USA to discuss this disease and create projects designed to expand research and improve treatment and prevention methods. The International Endotheliotropic Herpes Virus Workshop was sponsored by the Houston Zoo and the International Elephant Foundation (IEF), it brought together some of the world's foremost veterinarians, virologists, pathologists, researchers, biologists, and conservationists from Europe, Asia, Australia and North America.
Delegates to the workshop ended two days of discussions with the creation of eleven projects designed to expand research and improve treatment and prevention methods for EEHV. The projects in progress at the close of the workshop included establishment of a herpes reference lab in Asia, research to identify and evaluate cells and secretions which harbor the elephant herpes virus, a multifaceted study that will investigate alternative antiviral medications and determine treatment protocols based on the way the virus causes the disease, a retrospective study of pathology, disease triggers, and the effect of animal movements on the progression of the disease, and identification of sources of funding to continue research. Other projects identified were isolation and cultivation of the elephant herpes vaccine, investigation of cellular immunity, and identification of specific susceptible elephants. Another project in the initial stages is development of an Enzyme-Linked Immunosorbent Assay (ELISA) test to determine which elephants have been exposed to the virus.

Workshop delegates were in agreement that the two day session was extremely productive, and the goals, priorities, and timelines established at this workshop represent a significant step toward finding a cure for this deadly disease.



Artist: Manjula Wijesundara

Charlie Gray
Superintendent of Elephants, African Lion Safari,
Ontario,
Canada



Photo: Heidi Riddle for International Elephant Foundation (IEF)

Historical Notes

History and culture of elephants in Sri Lanka

Jayantha Jayewardene,

In Sri Lanka, no other animal has been associated for so long with the people, in their traditional and religious activities, as the elephant. This association dates back to the pre-Christian era, more than 5,000 years. Ancient Sinhalese kings captured and tamed elephants, which used to abound in the country. Various methods of capture were employed, some indigenous, others introduced by neighboring kings and countries that conquered and ruled Sri Lanka. Gradually the number of elephants captured increased. All elephants were kept by the king in his stables. The methods of capture were refined and modified, as time went on. Elephants, suitably caparisoned, have and still take part in ceremonial, cultural and religious pageants and processions.

Elephants have been used by man in his wars, in Europe and Asia. They have assisted him in his logging operations and construction works. In this country too, elephants have fought in wars and featured in various sports and combat, during Sinhala celebrations. In India they have provided transportation for sportsmen indulging in shikars. During the time of the Sinhala kings the elephant was afforded complete protection by royal decree. The penalty for killing an elephant was death. With the advent of the British, this protection was withdrawn. Large numbers of elephants were killed by the British under the guise of sport. Not only did the British government encourage and condone killings as a sport, but it also paid a bounty for each elephant killed, deeming the elephant an agricultural pest.

In Sri Lanka, the variations in physical appearance amongst elephants were noticed and recorded in ancient Sinhala manuscripts. There are ten such groups or 'castes'. These differences do not seem important now.

The first record of the association between man and elephant in Sri Lanka was recorded in the 1st Century BC, on an inscription at Navalar Kulam in Panama Pattu in the Eastern Province, of a religious benefaction by a prince who was designated Ath Arcaria or Master of the Elephant Establishment. The Elephant Establishment was called the Ath panthiya. The ruins of the ancient cities in Sri Lanka abound with carvings of elephants in many forms, attesting to the close association between man and elephant.

Jayantha Jayewardene

Biodiversity & Elephant Conservation Trust
615/32, Rajagiriya Gardens
Nawala Road, Rajagiriya
Sri Lanka

Sinhala literature of the 3rd Century BC indicates that the state elephant or Mangalahatti was the elephant on which the king rode. This elephant was always a tusker and had a special stable called the hatthisala. The post to which it was tethered was called the alheka. A 12th Century inscription on a stone seat at Polonnaruwa records that King Nissanka Malla sat upon it while watching elephant fights. These fights were staged for the entertainment of nobles. A rock sculpture of an elephant on the banks of the Mahaweli River was described thus, by archaeologist H.C.P. Bell: "This piece of animal sculpture is probably unique in Ceylon. Cut in full round from a rock, life-size, are the head and shoulders of an elephant whose feet the river washed when low. The elephant stands in the water, looking slightly upstream, as though hesitating to cross. At present the river in semi-flood reaches its eyes. There are signs of 'sets' for some building's foundations on a boulder adjoining, but no ruins or inscriptions are known likely to afford a clue to the object of this solitary tour de force of a skilful sculptor," (Bell & Bell, 1893). Unfortunately this rock sculpture no longer exists, having been blasted probably by fishermen dynamiting fish.

The first description of the capture of elephants in 40 AD is by Pliny. Here, the information that he gathered was from the Sinhalese ambassador to the court of the Emperor Claudius. Elephants were used on all important ceremonial occasions, especially where pomp and pageantry were required. The annual Perahera in Kandy, which dates back nearly 220 years, brings together well over a hundred elephants that parade the streets during the nights on certain pre-determined days in July-August each year.

New Year festivities in Sri Lanka feature elephants in various sports and competitive combat. Elephant fights were a popular form of Sinhala sport in early times and was called Gaja Keliya. Being built like a tank, elephants were used in war not only as a means of transport, but also as an instrument of defence and offence. They were used to ram barricades and, as Ivers, (1899) points out, "in time of war, they now and then fix a heavy iron chain to the end of their trunks, which they whirl around with such agility, as to make it impossible for an enemy to approach them at that time".

From the earliest of times there had been a significant demand for Sri Lankan elephants, from other countries. Aelian, (quoted by Emerson Tennent, 1859), says that the export of elephants from Ceylon to India had been going on without interruption from the

period of the First Punic War. India wanted them for use as war elephants, Myanmar as a tribute from ancient kings, and Egypt probably for both war and ceremonial occasions. The elephants from Sri Lanka were found to easily adapt to war, and were considered better than those from the mainland.

Their excellent qualities were well known to the Greeks even as far back as the 3rd Century BC, in the time of Alexander the Great. Onescritus, who was an Admiral of the Fleet of Alexander the Great and probably the first European to describe the trained elephants of Ceylon, has stated that the elephants from Taprobane (later Ceylon and then Sri Lanka) "are bigger, more fierce and furious for war service than those of India,". Greek writers like Megasthenes (circa 300 BC) and Aelian (44AD) corroborate this. Sixth Century writer Cosmos Indicopleustes says that the elephant from Sri Lanka was highly priced in India for its excellence in war. Elephants from Sri Lanka were exported to Kalinga by special boats, from about 200 BC, from the port of Mantai the present day Mannar. Such exports are also recorded by Ptolemy in 175 AD.

By this time Sri Lanka had also earned a reputation for skilled elephant management. The Sinhala kings had special elephant trainers. They were the Kuruwe people from Kegalle. Training elephants caught from the wild, for both traditional purposes and war, was the responsibility of these people. Even persons (mahouts) who looked after the elephants after their training, were trained by the Kuruwe people. A brass model of an elephant with a number of movable joints was used in the training of the mahouts.

Records show that even though Sri Lanka was exporting a large number of elephants in the 5th and 6th centuries BC, a number of elephants were also imported into the country after the 4th Century BC. This is apart from the gifts that the ruling monarchs of India and Myanmar, (then Burma) sent from time to time.

The Culavamsa (Ch. LXXVI) records that during the reign of King Parakrama Bahu (1153-1186 AD), King Ramana of Myanmar decreed that the practice of selling elephants from his kingdom for export should henceforth be stopped. "Moreover with evil intent, the king also set a high price on the beasts, commanding that the elephants which were sold in former times for a hundred nikkhalas of silver, or a thousand, should now be sold for two thousand or three thousand, and likewise he put an end to the ancient custom of giving an elephant to every ship that bore presents to the king [of Sri Lanka]."

The chronicle goes on to say that Prakrama Bahu made war on the King of Burma and subdued him. Later, the Burmese relented and said "Take henceforth from us as yearly tribute, as many elephants as are necessary." Records of the 12th Century AD again show that elephants continued, to be imported from Burma.

The export of elephants too continued and this is confirmed from time to time by writers on Ceylon - Sinhala Chronicles (15th Century), Athanasius Nikitin the Russian traveler (1470). Add-er-Razzak (1442) refers to the trade in elephants between Calicut and Ceylon.

Duarte Barbosa (1518) refer to the Royal monopoly of elephants - a good elephant fetched 1,500 ducats on the Malabar Coast at that time. Ribeiro (1847) states that "As the Ceylon elephant was superior, traders were prepared to pay twice or even up to four times for them, compared to elephants from other countries."

There are a number of references in early writings to man's association with and his use of elephants. The Mahawamsa (Sri Lanka's chronicle of history) details many such instances, especially that of Kandula, the elephant on which King Dutugamunu (200 BC), rode to war.

Dutch, Portuguese and British reports and books record several instances of elephant capture, their use by the Sinhala Kings in their armies, elephant fights and the execution of criminals by elephants. In certain instances the strength of a King or Potentate was judged by the number of elephants he used in war.

The King of Kandy maintained a special unit that dealt with all matters concerning elephants, including their capture, training, conservation and export. This unit was under the chief officer known as the Gajanayake Nilame. The Gajanayake Nilame, was of a high caste and received many favours, including land, from the king. The elephant catchers and keepers were from the lower castes.

During the times of the Sinhala kings, even though there were tens of thousands of elephants in all parts of the country, this animal was afforded complete protection by royal decree. Accordingly, no elephant could be captured, killed or maimed without the king's authority. All offenders were punished by death. Unlike today, the cultivators of that time could not plead that the elephants were harmed in the protection of their crops.

Any depredation or damage to crops by wild elephants had to be prevented by stout fencing, together with organized and effective watching by the farmers. It is interesting to note that though there were many more elephants then than now, Sri Lanka was considered to be the granary of the East.

When the Portuguese captured the maritime provinces of Sri Lanka they found a flourishing export trade in elephants. They too, quickly got involved in the elephant export trade, and at first obtained their elephants as tribute from the Sinhala people through their leaders. Thereafter they captured animals on their own. The Portuguese also set up a revenue-gathering unit, similar to the king's organization, known as the Elephant Hunt. Abeysinghe (1966) wrote that the Portuguese

maintained an annual demand of 37 elephants for export from two kraals. These were valued at 9,250 rix dollars which was equal to 15% of the total revenue of the state.

In 1507 the Viceroy of India sent a gift of a small elephant, imported from Ceylon, to King Manuel of Portugal. After seven years in Lisbon this elephant, named Annone, was presented to Pope Leo X and moved to Rome. Annone, lived in Rome for three years but died after developing stomach trouble due to the variety of food given to it by visitors and admirers. There is a memorial in Rome to Annone the first elephant in the Vatican (Hulugalle, 1969).

The King's Elephant Unit continued to operate within the Kandyan kingdom, even after the Portuguese occupation of the Maritime Provinces. Latterly however, the function of the King's Elephant Unit was only to supply the king's army with elephants. This was because with the development of cannons and musketry, the elephant was both frightened and vulnerable, and its export demand as an instrument of war was greatly reduced. During the reign of the Portuguese the person in charge of the Elephant Hunt was called the Gajanayake. The Gajanayake was in charge of stables at Matara. This was a large establishment. In 1697 there were 97 elephants in the stables at Matara. **Baldeus** wrote, in 1704, of a place in Matara, where captured wild elephants were tamed before they were sold to buyers who came from the Coramandel Coast and Bengal. A very large stable had been built to house these animals. These stables at Matara are the site of the present Kachcheri. The animals apparently were bathed twice a day in a nearby river, very likely the Nilwala. Tame elephants were used as monitors and trainers.

Those people who were sent into the jungles to look for suitable herds of elephants to be captured in kraals were called Baddenas. When the herds were sighted, the Dissawa of the area was informed, and he in turn gave instructions for arrangements to be made to hold the kraal. The men, numbering over a thousand, were divided into four groups under a leader called a Hattrebethmarale. The Aratchies were those in charge of the trappers who noosed the elephants once they were inside the Kraal and also trained the captured elephants.

In 1586, the king of Kandy, Rajasingha I, led an army which included a strong force of 2,200 highly trained elephants for fighting, and for other services, and laid siege on the Portuguese fort in Colombo. The siege however, was not successful. It is recorded that, in 1706, the king of Kandy had in his stables over 300 tuskers. The elephant was used less and less for war, and subsequently only for ceremonial occasions.

In 1656 the Dutch laid siege on the Fort of Colombo held by the Portuguese. Ribeiro, the Portuguese soldier and historian, records that all the elephants in the Fort excepting one, were eaten by the defenders as they ran short of food after a time. Only one elephant was spared, because it was needed to carry timber to repair the defences that were being damaged by the attackers.

Robert Knox, a Scotsman, who was a prisoner in the Kandyan kingdom for nearly 20 years, writing in the 17th century, stated "that the King makes use of them (elephants) as executioners: they will run their teeth [tusks] through the body, and then tear it in pieces, and throw it limb for limb. They have a sharp iron with a socket with three edges, which they put on their teeth at such times; for the elephants that are kept have all the ends of their teeth cut, to make them grow better, and they do grow out again."

Sirr (1850) also says that elephants were used as executioners of criminals, by training them to crush the victim's limbs, and by placing one of its legs on the man's body, tears off the limbs. Fortunately, the use of the mild tempered elephants for such gruesome executions has long been stopped.

Pybus (1762) states that the Dutch had to obtain permission from the king of Kandy to capture elephants which were within his domain. The king generally agreed to the Dutch capturing 20 to 30 animals each year, but the Dutch constantly exceeded this figure, capturing around 150 each year and 200 in one year. They continued to use the elephant stables at Matara referred to earlier. Elephants were also exported by the Dutch from Karativu island. The elephants were driven into the Jaffna peninsula by a shallow ford that separated it from the mainland. This ford has now been bridged and given the name Elephant Pass.

The Dutch held an annual sale of elephants in Jaffna. Elephants caught in Kraals and those received as tribute were sold there. Buyers from the Coramandel and the Bengal coasts continued to attend these sales regularly.

The traditional methods of capture by noosing etc. were practiced for a much longer period. However, in 1761 the Dutch Governor Becker made a decree prohibiting the use of pits and nooses for the capture of elephants. This was in a bid to keep the mortality rate among the captive elephants low. The form of noosing practiced in northern Ceylon was different to the noosing methods of the Eastern Province and other parts of the country. A large noose was suspended from a strong tree, with a man or several men on the tree to manipulate the noose. Elephants were then driven towards the tree with the noose. With this ban, kraaling became the only method of capture that could be employed.

When the British captured the Maritime Provinces from the Dutch in 1796, and later the Kandyan Kingdom in 1815, they continued the capture of elephants for some time, but on a low-priority basis. The British however, indulged in the shooting of elephants as a form of sport. Elephant populations that had been able to withstand the detrimental effects of capture all these years, now started diminishing rapidly with the wanton and indiscriminate

destruction of the elephant herds. Major Thomas Rogers is credited with having shot over 1,500 elephants. This works out to an average of one elephant being killed by him every day for four years. Two others, Captain Gallwey and Major Skinner are reputed to have shot half that number each. Many other 'sportsmen' have shot in the region of 250-300 animals during this time.

As the elephant was a threat to the agricultural activities of the rural population, the British provided guns freely to villagers to keep away the marauding elephants from their cultivations. This action, which seemed necessary at that time, added to the destruction of the elephant. Farmers, who had hitherto protected their crops from marauding elephants by other means, now had a much easier method. They shot at them and either maimed or killed them.

The British were also interested in developing plantation crops in addition to subsistence crops. British planters, who were opening up the railways and roads along with coffee and later tea plantations, also shot trespassing elephants at will. Here again, the purported protection of their crops seemed to justify their actions. The planters combined their sport and the protection of their plantations and shot elephants at will, so much so that the once large elephant population in the hills dwindled rapidly.

The British did away with the Elephant Department started by the Sinhalese, and the Elephant Hunt maintained for the occupied areas by the Portuguese and the Dutch. They also greatly reduced the number of kraals that were held. In fact in 1828 the British passed a law prohibiting the capture of elephants except for the government. This law was rescinded in 1831. The Kandyan chiefs, however, continued holding kraals, and it is recorded that from 1800 to 1900, fifty two kraals were held. The last Kraal was held in 1952.

References

Abeysinghe, T. (1966). *Portuguese Rule in Ceylon, 1594 -1612*, Lakehouse Bookshop, Colombo

Baldeus P. (1704). *Description of the great and most useful Isle of Ceylon*, New Delhi, 1996

Barbosa, D.(1518). *An account of the countries bordering on the Indian Ocean and their inhabitants,* (ed. M.L. Dames), London

Bell, B.N. & Bell H.M. (1893). *H.C.P. Bell Archaeologist of Ceylon and the Maldives*, Archetype Publications, Clwyd.

Hulugalle, H.A.J. (1969). *Ceylon of the early Travelers.* Colombo.

Ivers, R.W. (1899). *Manual of the North Central Province of Ceylon.* Government Printer, Ceylon

Pybus (1762). *The Pybus Embassy to Kandy 1762.* R. Raven Hart; (ed. P.E.P. Deraniyagala) Colombo 1958

Ribeiro, J. (1847). *History of Ceylon, presented by Captain John Ribeiro to the King of Portugal in 1685* translated by George Lee, Colombo

Sirr, H.C. (1850). *Ceylon and the Cingalese.* London



Photo: courtesy Dr. H. I. E. Katugaha

Book Review

Endangered Elephants: Past, Present and Future

All over their range elephants come in close contact and conflict with people. The elephant, with its massive bulk and power, social organization, behaviour, and intelligence, has been a figure of reverence in religion, a cultural symbol, a useful work animal that has fascinated man from the earliest of civilization. While this is true of all elephant-containing geographical areas of the world, in Asia especially, people have nurtured close ties with the elephant for centuries. In Asia also, are the worst incidences of human-elephant conflict when both man and beast turn against each other resulting in fatal clashes that claim lives on both sides.

Many of the elephant habitats of Asia are densely populated. The agricultural practices in many of these countries encourage elephants to move into human habitation. The elephant's great adaptability to changed environments and its preferences to open, forest edges places the animal in more danger of direct contact with humans.

But of the world's elephant population, the majority of the animals live in Africa. Although there are very large natural reserves and less population density, human-elephant conflict is not rare in Africa. The root causes of the conflict and the socio-economic situation of the people involved are startlingly similar in both Asia and Africa.

In 2003, Colombo (Sri Lanka) hosted a landmark symposium that brought researchers and conservationists from 23 different countries of Asia and Africa to share lessons between the two vastly differing continents and come up with a way forward that would ensure the survival of the elephant species. The three-day symposium was titled 'Human Elephant Relationships and Conflicts' and served to expose new conservation techniques, unveil new and exciting elephant research and present many past experiences of dealing with conflict between man and elephant.

The legendary Iain Douglas-Hamilton from Africa in his opening address to the forum spoke emotionally about the plight of elephants all over the world. "My heart bleeds when I read the endless and almost hopeless catalogue....', he said. "Elephants are weekly being shot, snared, electrocuted, run into trains, poisoned and everywhere deprived of habitat."

Douglas-Hamilton posed the question – do we leave elephants all alone to the wilds and hope they survive the best they can, or as certain conservationists insist, actively manage wildlife so that that species have better chance of survival especially if they have an economic value?

The symposium resulted in over 50 papers on a wide range of elephant-related topics presented by scientists, field researchers, funding agencies, zoological gardens, veterinary surgeons and conservationists. The topics ranged from historical perspectives of human elephant relationships to genetic research, to documentation of conflict in different countries and ranges, management of domesticated elephants and traditional knowledge in elephant capture. These papers are now published in a book titled **"Endangered Elephants: Past, Present and Future."** By publishing these papers with their accompanying maps, photos was primarily, to fulfill the aim of making this information available to a wider audience than the 220 delegates who participated at the symposium. The research, the field experience and the methodologies discussed in the book under different subject headings would lead to better understanding of the species and their conservation needs in areas that are becoming highly populated and where agriculture is taking over previous elephant habitat.

Jayantha Jayawardene, editor of the volume, who was instrumental in organizing the symposium said, "This book will be a substantial addition to the scientific information out there on the elephant and contribute greatly to future conservation practices. It is a great resource for practitioners and researchers alike, since rarely have we managed to get experiences of such a wide range of countries in to a single volume."

The final discussions at the 2003 symposium led to an agreement that the forum would request the United Nations to establish an inter-governmental body with scientific support from the IUCN Specialist Groups on Elephant Conservation. The aim of such a body would be to coordinate initiatives within a global strategy for future survival of the species and also to encourage inter-agency communication and facilitate synergies. The Body would also be able to assist range states in the development and implementation of policies that would protect elephant ranges. This letter, signed by Debbie Olsen of International Elephant Foundation (USA) and Jayantha Jayawardene of Biodiversity and Elephant Conservation Trust (Sri Lanka) is also included in the book.




Photo: courtesy Richard Lair (Thailand)

Miscellaneous

IUCN/SSC Asian Elephant Specialist Group - Announcement of new Co-Chairs

Dear colleagues,

The Chair of the IUCN Species Survival Commission (SSC) has invited us to serve as interim Co-Chairs of the Asian Elephant Specialist Group (AsESG), and we have accepted the invitation. This means that the Group will not have a single Chair as before - instead we will share the Chair's responsibilities. The IUCN process of appointments is a formal and time-consuming one; with all Chairs and members being appointed or reappointed in the new quadrennium. Fortunately, the Specialist Group Chair invitations are now complete, and so we can begin the process of (re)appointing the AsESG's members.

A major task for the AsESG is to help mould and guide decisions and policies affecting elephant conservation by providing sound scientific information and advice. The SSC has charged us with delivering the following objectives over the coming year:

1. Establish a new, revitalized AsESG membership;

2. Establish a regular dialogue with the new membership including timely updates in Species, SSC's E-Bulletin, or through a dedicated AsESG publication or website;

3. Provide support and motivation for the members of the AsESG to stimulate their technical input and activity to new and ongoing activities in support of the conservation of Asian elephants, including providing technical support to the upcoming Asian Elephant Range States meeting;

4. Revise and complete an updated version of the Asian Elephant Action Plan;

5. Achieve agreement within the AsESG on plans and priority actions for the Group;

6. Help identify sources of long-term funding support for the core functions of the AsESG;

7. Other actions deemed necessary to build the capacity of the AsESG and its members in South and South East Asia.

We see working with the range State governments to effect better conservation for Asian elephants as a particularly high priority for the group. Here we recognize that the group's advice would be most acceptable if it originates from and has the neutrality of an IUCN Specialist Group behind it. This necessitates that the group be a purely neutral body whose members do not have any vested interests driving their decision-making. The group will also focus on providing technical support and practical solutions to conservation problems and on increasing the range States' capacities to manage their elephant populations.

We recognize that the Asian elephant situation is precarious in many ways and needs more global/regional attention and support. One way we will be working to achieve this will be by providing a clear global perspective on the status of the Asian elephant, the threats faced by it, and the conservation issues involved. We believe the AsESG should create a 'database' on the status of Asian elephant populations and habitats and link this to a 'strategic plan' that identifies and addresses specific needs and priorities within individual countries. The AsESG will endeavor to provide governments, NGOs, donors and individuals involved in Asian elephant conservation with guidance on how best they can work and contribute at the local, national, regional, and global levels. Furthermore, we recognize that setting research and conservation priorities, funding, institutional capacity, legislation, and captive elephant management are among the many important issues that need the group's attention, but the details of how the AsESG will address these issues are for the group's members working with the Co-Chairs to decide and work on. Achieving the above will be possible only through a membership that is active, representative, and includes managers and scientists alike. Communication will be a priority as the group cannot function effectively without establishing reliable channels of communication. In building the group's membership we will pay attention to the lessons learnt by the IUCN/SSC's other specialist groups. For example, "While long experience has shown that smaller groups tend to be more effective, lasting conservation action in the international sphere almost requires that Groups be more than small coteries of like-minded colleagues." We do realize the group cannot be so large as to include everyone interested in or working with Asian elephants and we have been given important advice on the appointment of members and composition of SGs from the SSC, but we would like everyone to know that we do not look at the AsESG as 'the only' interested party. Others who have technical expertise will be invited to work and contribute through 'Task Forces' that will be constituted to deal with specific technical issues. Additionally, where issues are of a more general nature and require wider consensus we will seek the opinions/views of the wider Asian elephant conservation community before presenting an AsESG perspective on the most important issues.

We look forward to working with you to achieve these aims and to ensure a better future for Asia's elephants.

Sincerely,

Ajay Desai & Simon Hedges

Interim Co-chairs,
IUCN/SSC Asian Elephant Specialist Group




Photo: Heidi Riddle for International Elephant Foundation (IEF)

Miscellaneous

Ringling Bros. and Barnum & Bailey Commits $180,000 to Smithsonian's National Zoo to Continue Critical Elephant Herpes Research

Ringling Bros. Funding Will Support Search for Cure to Single Largest Threat to the Endangered Asian Elephant

Vienna, VA, (July 25, 2005) -- *Ringling Bros. and Barnum & Bailey*®, today announced it will fund ongoing research on endotheliotropic herpes viruses (EEHV) being conducted by the National Elephant Herpesvirus Laboratory at the Smithsonian Institution's National Zoological Park. The $180,000 gift will be distributed over the next three years to support the National Zoo's efforts toward treating, and ultimately curing, this typically fatal disease in young Asian elephants. The gift reflects *Ringling Bros.*' commitment to the survival and well-being of the endangered Asian elephant.

EEHV is the single greatest health threat to the Asian elephant today. Only four known elephants have survived the disease, one of which is a *Ringling Bros.* elephant currently residing at the *Ringling Bros. and Barnum & Bailey Center for Elephant Conservation*, a 200-acre facility in central Florida dedicated to the reproduction, study and retirement of Asian elephants. The National Zoo's EEHV lab is the only facility in North America that tests for the disease. EEHV infects only elephants and usually has a fatal outcome within a week of onset of symptoms, which include head and neck swelling, tongue cyanosis and lethargy. Elephants with EEHV symptoms are immediately started on famciclovir (an anti-herpes drug used in humans) to increase their chance of survival.

"As a partner in the fight to preserve the endangered Asian elephant, we applaud the National Zoo for its work in protecting this magnificent species from this deadly virus," said Bruce Read, Vice President of Animal Stewardship for *Ringling Bros.* "We hope that our donation will ensure the continued existence of the zoo's EEHV lab and the critical research and testing that will someday lead to the eradication of this disease."

In 1995, pathologists at the National Zoo were the first to identify EEHV while investigating the sudden death of a 16-month-old Asian elephant born at the zoo. The zoo's EEHV lab tests blood samples from symptomatic elephants to confirm they are suffering from the disease. Additionally, the lab is currently developing a test called Enzyme-Linked Immunosorbent Assay (ELISA) that will help identify at risk elephants.

"The research and services provided by the National Elephant Herpesvirus Laboratory at the Smithsonian's National Zoo are a critical component to the collective efforts of the elephant community for the preservation of these endangered species," said Dr. Laura Richman, former pathology resident and National Zoo research associate, who founded the EEHV lab. "It is our hope that the lab will generate meaningful diagnostic and epidemiologic data that will help curtail illness and deaths attributable to the virus."

Young Asian elephants are especially vulnerable to EEHV, as about half the deaths of young elephants in captivity are attributed to the disease. The only clear risk factor currently identified for Asian elephants is exposure to African elephants, although infection is not limited to this factor.

"There is more research that needs to be done in order to develop effective medical treatments for EEHV," said Mike Keele, Chair of the American Zoo and Aquarium Association Elephant Species Survival Program and Deputy Director of the Oregon Zoo. "Unlike many groups that claim they care deeply about elephants, *Ringling Bros.* is making a major resource commitment to improve their lives while in our care. *Ringling Bros.*' generosity is an example of cooperative efforts that can truly make a difference for elephants worldwide."






Photo: Heidi Riddle for International Elephant Foundation (IEF)

Miscellaneous

International Elephant Foundation
2007 Elephant Conservation and Research Funding Support

The International Elephant Foundation (IEF) is accepting proposals for elephant conservation and research funding support.

IEF is offering financial support in 2007 for *in situ* and *ex situ* projects, including protection of wild elephants and their habitats, scientific research, education efforts, and improvements in captive elephant care. Proposals are peer-reviewed by a panel of advisors from the fields of medicine, research, academia, and elephant management. **Please read the instructions closely. Proposals must follow the format outlined below. Those applications that do not follow the instructions will not be considered for funding.**

Awards typically range about US$10,000 per year.

Proposals are preferred that meet some or all of the following objectives and criteria:

- The proposal should clearly contribute to the *in situ* or *ex situ* conservation of African or Asian elephants or their habitats.
- Project is part of an established conservation program or is well-suited to become a long term program.
- Project has conservation value and measurable impact.
- Project is grounded in sound scientific methodology, is logistically feasible, and has a high probability of success.
- Project is collaborative in nature.
- Project has multi-institutional participation and matching funds.
- Project and Principle Investigator demonstrate a spirit of cooperation with *ex situ* elephant facilities and other like-minded conservation institutions.

Examples of some funding priorities are:

- Capacity building
- Strategies for human elephant conflict resolution
- Strategies to counteract the bushmeat crisis/ ivory poaching
- Strategies to manage local elephant over-population problems
- Translocation follow-up studies
- Ex Situ elephant management, veterinary and reproduction projects

Proposals must be in English and budgets in US currency.

Proposals must be received at the IEF office by the end of the day on 1 September 2006.

Proposals must be submitted electronically as ONE attachment, this includes the cover page, narrative, CVs, supporting documents etc. Photos are not permitted unless they are VITAL to the clarification of the proposal. In that case, photos must not be imbedded in the application but sent as separate attachments.

Proposals must be submitted electronically to dolson@indyzoo.com.

Frequently Asked Questions

When is the application due?
The deadline is September 1, 2006. Late proposals cannot be considered out of fairness to others.

Can I submit more than one proposal?
You may submit two or more proposals, recognizing that the total number of proposals submitted does not necessarily enhance your ability to be funded.

Who is eligible?
We encourage global projects. Organizations outside the US are eligible as are students, scientists and institutions. Students are required to file applications through a faculty advisor; the advisor co-signs the applications and accepts reporting responsibility. Proposals should be submitted in English and in US dollars.

What about overhead costs?
Overhead costs are not funded.

What size grants do you entertain?
Requests of $10,000 or less are preferred but larger amounts will be considered.

Do you support multi-year proposals?
Two or three-year proposals will be considered but funding will only be awarded for 2007 with an application for each additional year required for multi-year proposals with no guarantee of funding. Reports will be required prior to award and distribution of funds for subsequent years.

How long is the process?
The review process is completed December 15, 2006.

Please do not call any earlier regarding the status of your proposal. It is our policy to notify all applicants at the same time regarding the status of their application. If you have not heard from us by January 31, 2007, please feel free to contact us then.

If I am funded, when are reports due?
Upon signature on the IEF 2007 Grant Program Contract, the 1st installment of funds will be made available to applicant. A six-month interim report, including digital photos, is required prior to receipt of the second installment. A final report and final budget, including digital photos, is required at the end of the funding year. Any future requests for funding will not be given consideration unless the six-month interim report and digital photos have been submitted, as well as all other contractual requirements. New applications from the same organization or individual will not be considered if reports from funded projects in prior years have not been received.

INTERNATIONAL ELEPHANT FOUNDATION APPLICATION FOR IEF FINANCIAL SUPPORT
Cover Sheet

TITLE OF THE PROJECT:

Principle Investigator
Name & Title:
Institution:
Institution Address:
Phone, Fax, email:

Co-investigator(s)
Name & Title:
Institution:
Institution Address:
Phone, Fax, email:

Name & Title:
Institution:
Institution Address:
Phone, Fax, email:

Budget
Total Project Budget:
Total Requested from the International Elephant Foundation:

Project Start Date:
Project End Date:
Is the Project Underway?

Project Category
Zoo Research
Field Research
Field Conservation
Conservation Education
Professional Training/Technology Transfer/Capacity Building
Habitat Protection
Applications must include the following:

1) **COVER SHEET** (sample attached above) containing project title, name and contact information for the principal investigator, name and institutional affiliation of co-investigator(s), amount requested, project start and end dates, and project review category.

2) **ONE-PAGE SUMMARY** containing;

   - summary of the project and project objectives
   - anticipated benefits and outputs
   - personnel & organizations involved

3) **PROJECT NARRATIVE** of a maximum of 10 pages, 10-12 point font, containing:

   - project rationale
   - goals & objectives
   - methodology & work plan
   - feasibility
   - monitoring & evaluation procedures
   - sustainability
   - potential significance of results to conservation

4) **TIME LINE** with proposed Start and End dates. Also Please note if the project is currently underway.

5) **BUDGET** must be itemized and divided into the following categories: project personnel (stipends or salaries support, number of persons, hours, rate, position), travel expenses (number of persons, cost per trip, number of trips), lodging and meals (numbers of days x persons), equipment, supplies, services, miscellaneous expenses. Please also list funds from other sources that are supporting the project. The following sample budget format must be used:

BUDGET ITEM	AMOUNT REQUESTED FROM IEF	MATCHING FUNDS	SECURED FROM OTHER PARTNERS	BALANCE REQUIRED	TOTAL PROJECT COSTS
Field Patrol per diem 2 patrols/wk x 21 staff @ $10 ea					
Elephant subsistence 10 elephants x 12 mo @$100/mo					
CRU unit uniforms for 30 staff @$20 ea					
TOTALS					

1) Please justify the budget requested. Please also address what will happen/what your alternatives will be foar this project if it is not chosen or awarded the full amount you are requesting for funding from IEF.
2) Provide a map showing the location of your project site at the state or province level.
3) **QUALIFICATIONS** of principle and co-investigators (**maximum 2-page CV per individual**)
4) Indicate whether **PERMITS** or **IACUC** approval (where required for the study) have been obtained.

Applications must be in English and received at the IEF office by the end of the day on 1 September 2006.
Applications must be submitted electronically to dolson@indyzoo.com.

For more information and questions contact:

Deborah Olson
Executive Director

International Elephant Foundation

P.O. Box 366
Fax: 817-237-7183
Azle, Texas 76098
E-mail: dolson@indyzoo.com

Instructions for Contributors

GAJAH is the official journal of the Asian Elephant Specialist Group (AsESG) of the Species Survival Commission (SSC) of the World Conservation Union (IUCN). The journal is intended as a medium for communication and discussion by the members of the AsESG of important issues that concern the conservation and management of the Asian Elephant (Elephas maximus) both in the wild and in captivity. GAJAH welcomes communications and research papers on all aspects of the Asian elephant with the focus on its conservation and management. GAJAH is aimed at professionals, biologists and academics carrying out research on Asian elephant, government and non-government organizations involved in its conservation, and interested members of the general public. All articles published in GAJAH, are deemed to reflect the individual views of the authors and not the official points of view, either of the Asian Elephant Specialist Group (AsESG) or the Species Survival Commission (SSC). GAJAH is a non-profit publication that is supported by financial assistance from the U.S. Fish and Wildlife Service.

Categories of manuscripts

Research articles report research results, and should include a short abstract, introductory paragraph, brief subheads, and references.

Short communications must contain important new findings that are of general interest.

Notes from the field: GAJAH welcomes notes from the field.

Historical notes concern interesting aspects of personalities and events in elephant conservation and management

Miscellaneous includes letters, book reviews and announcements of forthcoming meetings related to elephant conservation.

Preparation of manuscripts

Manuscripts should be in British English. They should be typed double-spaced on A4 paper. Electronic submission (in MS Word) is also acceptable. The manuscript should have an abstract. References should be cited in the text as, for example, Eisenberg & Lockhart (1972) or (Sukumar, 1989). For three or more authors, use the first author's surname followed by *et al.* Multiple references should be in chronological order. The reference list should be in alphabetical order following the Harvard author-date system. Write out journal titles in full. Acknowledgements should be as brief as possible. In the reference list, cite the publications as follows.

Sukumar, R. (1989) *The Asian Elephant: ecology and management.* Cambridge Studies in Applied Ecology and Resource Management. Cambridge University Press, Cambridge

Barnes, R.F.W. (2001) How reliable are dung counts for estimating elephant numbers? *African Journal of Ecology,* **39:** 1-9

Kahl, M.P. & Armstrong, B.D. (2000) Visual displays of wild African elephants during musth. *Mammalia,* **66:** 159-171

Hanks, J. (1981) Characterization of population condition. In *Dynamics of Large Mammal Populations* (eds. C.W.Fowler & T.D.Smith), pp. 47-73. John Wiley & Sons, New York.

Olivier, R.C.D. (1978) On the ecology of the Asian elephant *Elephas maximus* Linn. with particular reference to Malaya and Sri Lanka. Unpublished Ph.D thesis, University of Cambridge, U.K.

All manuscripts must be addressed to Dr. Charles Santiapillai, Editor, GAJAH, Department of Zoology, University of Peradeniya, Sri Lanka. e-mail: csantiapillai@gmail.com

